

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 : C07D 475/02, 473/04, 513/04, 239/96, 217/24, 401/14, A61K 31/505, 31/52 // (C07D 513/04, 285:00, 239:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/00491 (43) International Publication Date: 6 January 2000 (06.01.00)</p>
<p>(21) International Application Number: PCT/US99/14320 (22) International Filing Date: 24 June 1999 (24.06.99) (30) Priority Data: 09/107,026 29 June 1998 (29.06.98) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). (72) Inventors: CARSON, Dennis, A.; 1482 Vista Del Oceano, Del Mar, CA 92014 (US). COTTOM, Howard, B.; 1890 Lenpee Drive, Escondido, CA 92025 (US). CHAO, Qi; Apartment 32, 4158 Decoro Street, San Diego, CA 92122 (US). (74) Agent: BERLINER, Robert; Fulbright & Jaworski L.L.P., 29th Floor, 865 South Figueroa Street, Los Angeles, CA 90017-2576 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
<p>(54) Title: BIFUNCTIONAL ANTAGONISTS OF CYTOKINE-SENSITIVE PROTEIN KINASE ACTIVATION CASCADES AND METHODS FOR USE AS ANTI-INFLAMMATORY AGENTS</p> <p>(57) Abstract</p> <p>Bifunctional activation cascade antagonist compounds inhibit inflammatory responses associated with TNF-α and fibroblast proliferation <i>in vivo</i> and <i>in vitro</i> by blocking activation of both the c-Jun and p38 kinases. The compounds of the invention possess greater than additive activity as compared to the individual constituents of each compound. One such constituent blocks activation of c-Jun kinase while the other blocks activation of p38 kinase, in a non-overlapping manner. The compounds of the invention neither appreciably inhibit the activity of cAMP phosphodiesterase nor the hydrolysis of phosphatidic acid, and are neither cytotoxic nor cytostatic. Methods for the use of the novel compounds as anti-inflammatory agents are also described. The methods are expected to be of use in reducing or protecting a vertebrate host against inflammatory responses (for example, after angioplasty), in limiting fibrosis (for example, of the liver in cirrhosis), in inhibiting cell senescence, cell apoptosis and UV induced cutaneous immune suppression.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PATENT

5 BIFUNCTIONAL ANTAGONISTS
OF CYTOKINE-SENSITIVE PROTEIN KINASE ACTIVATION CASCADES
AND METHODS FOR USE AS ANTI-INFLAMMATORY AGENTS

RELATED PATENT APPLICATIONS

 This application is a continuation-in-part of US Patent Application Serial
Number 08/858,778 filed May 19, 1997 (now pending), which is in turn a
continuation-in-part of US Patent Application Serial Number 08/367,102, filed
10 December 29, 1994 (now pending), which is in turn a continuation-in-part of US
Patent Application Serial No. 08/482,551, filed June 7, 1995 (now allowed).

STATEMENT OF GOVERNMENT SUPPORT

 This invention was made with government support under Grant No. GM-
23200 awarded by the National Institutes of Health. The government may have
15 certain rights in the invention.

BACKGROUND FOR THE INVENTION

1. *Field of the Invention*

 The invention relates to antagonists of cytokine-sensitive, mitogen-
activated protein kinase activation and their use as anti-inflammatory agents.

2. *History of the Related Art*

Two of the MAPK (mitogen activated protein kinase) signalling cascades are initiated by proinflammatory cytokines (PICs) and other stress factors, such as ultraviolet light, ionizing radiation, tissue trauma, cytotoxic agents (e.g., anisomycin and arsentine), heat shock and ceramide. These "PIC sensitive" signalling cascades lead to activation of the c-Jun N-terminal protein kinase (JNK; also known as SAPK, for "stress-activated protein kinase") and a newly discovered protein kinase, p38 (also known as CSBP, for "cytokine suppressive anti-inflammatory drug binding protein").

Activation of JNK and p38 is related to the onset and maintenance of acute and chronic inflammatory conditions such as localized fibrosis, cystic fibrosis, ultraviolet light induced cutaneous immune suppression, cell senescence and apoptosis, gastrointestinal inflammation (e.g., inflammatory bowel disease) and pulmonary inflammation (e.g., chronic bronchitis and asthma). Inhibition of PIC synthesis and/or release by stressed cells can reduce MAPK activation-related inflammation by avoiding initiation of the PIC sensitive MAPK signalling cascades.

Different anti-inflammatory agents approach this goal in different ways. For example, pentoxifylline suppresses synthesis of PICs by inhibiting cAMP phosphodiesterase activity. Corticosteroids inhibit transcription of PICs. Cytokine suppressive anti-inflammatory drugs (CSAIDs) are believed to inhibit PIC synthesis by binding a MAPK required for cytokine mRNA translation; these compounds selectively interfere with activation of the p38 signaling cascade. None of the drugs are completely effective in suppressing PIC synthesis or release and many have side-effects that are use-limiting.

SUMMARY OF THE INVENTION

The invention provides compounds which are bifunctional in that they block PIC initiation of both of the PIC sensitive MAPK activation cascades. In particular, the invention combines a novel antagonist of the JNK activation cascade with another PIC inhibitor, most preferably an antagonist of the p38 activation cascade. The components of the resulting "bifunctional activation cascade

antagonists" act synergistically to antagonize PIC-sensitive MAPK activation to a greater extent than could be achieved by either compound alone.


In particular, the bifunctional activation cascade antagonists of the invention inhibit up to 100% of PIC release by stressed cells. In contrast, 70-80%
5 PIC inhibition is the maximum level achievable by known PIC-sensitive MAPK antagonists. Advantageously, bifunctional activation cascade antagonists do not pose the risk of side effects associated with certain anti-inflammatory agents, such as the sleeplessness and anxiety induced by methylxanthine-based cAMP phosphodiesterase inhibitors (e.g., pentoxifylline). Further, the potency of the inventive compounds
10 permits their use at lower dosages than are required by use of their constituents alone.

The JNK activation cascade antagonist constituent of the inventive bifunctional activation cascade antagonists consists of heterocyclic molecules with biologically active side chains; specifically, purine, pteridine, thiadiazolopyrimidine, quinalozine and isoquinolone based compounds and water-soluble morpholinoethyl
15 esters thereof. Such compounds and methods for their synthesis are described in detail in commonly owned US Patent Application Serial Nos. 08/858,778, 08/367,102 and 08/482,551, the disclosures of which are incorporated herein as though set forth in full. The p38 activation cascade antagonist constituents are any compound which interferes with the initiation or completion of the p38 activation cascade. In
20 one aspect of the invention, the p38 activation cascade antagonist is a pyridylimidazole. An especially useful class of pyridylimidazoles for use in the invention are the CSAIDs.

The JNK and p38 activation cascade antagonist constituents of the bifunctional activation cascade antagonists are preferably conjugated together. Thus,
25 in one aspect of the invention, the constituents of the bifunctional activation cascade antagonists are conjugated to one another via a bond which is severable *in vivo*, such as ester, amide or azo linkages. In another aspect of the invention, unconjugated JNK and p38 activation cascade antagonist constituents are mixed in a pharmaceutically acceptable carrier. In yet another aspect of the invention, conjugated or unconjugated

JNK and p38 constituents of the bifunctional activation cascade antagonists are combined with a delivery vehicle, such as a colloidal dispersion system.

The invention further encompasses the following conjugated bifunctional activation cascade antagonists:



5 wherein n is any number of carbon atoms from 1 to 7, O or N;
R₁, if present, is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or
aralkyl having less than 7 carbon atoms;
R₆ and R₇ are H, OH or OR₁, in any combination;
Z is N or C;
0 X, where Z is C, is H, halogen, N₃, NO, NH₂, NHR₁, N(R₁)₂ or COR₁;
and,
A is H, halogen, N₃, NO, NH₂, NHR₁, N(R₁)₂ or COR₁.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 is a bar graph depicting inhibition of cell growth arrest in 3T3 fibroblasts according to the invention after growth arrest was induced through deprivation of the cells of serum. The cells were incubated and grown to 90% confluence in serum. The medium was then removed and replaced with serum-free medium. To assess the effect of an inventive compound (no. 37, a pteridine) on cell senescence in the presence of ceramide, aliquots of the cells were incubated with different concentrations of each. Concentrations of compound no. 37 are indicated by the insert legend while concentrations of ceramide are indicated along the x axis. Inhibitory effects were assessed as a measure of DNA synthesis; [^3H] thymidine incorporation detection is indicated along the y axis.

FIGURE 2 is a bar graph depicting inhibition of cell apoptosis in human (Jurkat) T lymphocytes according to the invention. The inhibitory activity of two inventive compounds (nos. 37 and 6 (a purine)) was tested in comparison to like activity of pentoxifylline and a control compound, Ro 20-1724. Activation of the sphingomyelin signal transduction pathway was stimulated by incubation of the cells with an anti-FAS monoclonal antibody (which binds CD95, a cell surface receptor which triggers cell apoptosis). Percent inhibition was measured as a function of the number of cells which excluded vital dye erythrosin B. Percent inhibition is indicated along the y axis while the concentration of compounds tested is indicated along the x axis.

FIGURE 3 is a bar graph depicting inhibition of activity on the part of CaPK in Jurkat cells according to the invention. The inhibitory activity of a compound of the invention (no. 37) was tested in the presence of either ceramide or anti-FAS. Inhibition of CaPK activity was measured as a function of phosphorylation and detected by autoradiography. The compounds the cells were incubated in are indicated along the y axis while the percent control (i.e., inhibition of CaPK) is indicated along the x axis. Shorter bars indicate greater relative inhibition.

FIGURES 4 (a) through (g) are copies of spectrographs indicative of absorbance of inventive compounds no. 37, no. 6, no. 37 in combination with no. 6, oxo variants of nos. 37 and 6, as well as, for comparison, PABA (p-amino benzoic acid, a common sunscreen additive) and isoquinolone. The inventive compounds
5 absorbed through most of the UVB wavelength, while a mixture of compound nos. 37 and 6 absorbed throughout the UVB wavelength.

FIGURES 5(a) and (b) depict, respectively, the results of an enzyme-linked immunosorbent assay (ELISA) for TNF- α production by bacterial lipopolysaccharide (endotoxin) stimulated human monocytes incubated with the
10 compounds of the invention and a control compound (Ro-1724, that is a known and specific inhibitor of phosphodiesterase type IV [the predominant isoform of phosphodiesterase found in monocytes and neutrophils]). Compounds tested are identified by the number assigned to them in Table 1. The horizontal axis of each graph shows the amount of each compound tested (in μ M) while the vertical axis
15 shows the IC₅₀ values for TNF- α production as a percentage of the production in the presence of only the control compound.

FIGURE 6 is a graph depicting the results of an assay for in vivo leukopenia in mouse blood in response to lipopolysaccharide (LPS). Leukopenia induced by LPS is mediated by TNF. Hence, this model assesses both TNF
20 production and action. Compounds tested for inhibition of leukopenia are identified by the number assigned to them in Table 1. Along the x axis of the graph, the numbers correspond to the number of white blood cells detected as cells/ml of fluid.

The results (shown by bars) are expressed in terms of a percentage of the leukopenia response (based on neutrophil content) to pure LPS, in absence of other compounds.

FIGURE 7 depicts the results of an assay for inhibition by compounds of the invention (nos. 37 and 6) of the effects of a cell permeable ceramide analog (C_2 - ceramide), dihydro ceramide and diacyl glycerol on TNF- α production by human monocytes. Inhibition of TNF- α production was measured by ELISA; the results are indicated in pg/ml of TNF- α along the x axis.

FIGURE 8 depicts the results of an assay for inhibition by a compound of the invention (no. 37) to prevent the stimulatory effects of C_2 - ceramide or protein kinase C activity in human lymphocyte extracts. Inhibitory effects were assessed as a measure of DNA synthesis; [3H] thymidine incorporation detection is indicated along the y axis.

FIGURE 9 depicts the results of an assay for *in vitro* TNF- α production by human macrophages in response to lipopolysaccharide (LPS) and inhibition of that production by pteridine and isoquinolone compounds of the invention (nos. 37 and 11-49). Along the x axis of the graph, the numbers correspond to the concentration of TNF- α detected in pg/ml.

FIGURE 10 depicts inhibition of PDGF induced fibroblast proliferation among 3T3 fibroblasts in response to the inventive compounds. The compounds tested are identified along the x axis by the numbers assigned to them in Table 1. Inhibitory effects were assessed as a measure of DNA synthesis; [3H] thymidine incorporation detection is indicated along the y axis.

FIGURE 11 depicts inhibition of EGF induced fibroblast proliferation among 3T3 fibroblasts in response to the inventive compounds. The compounds tested are identified along the x axis by the numbers assigned to them in Table 1. Inhibitory effects were assessed as a measure of DNA synthesis; [³H] thymidine incorporation detection is indicated along the y axis.

FIGURE 12 depicts data indicative of the apoptotic protective characteristics of the compounds of the invention as represented by compounds 1C-261 (Compound 37) and 1I-49. Human lymphocytes were cultured in serum aliquots with the concentrations of the inventive compounds indicated along the x axis of the **FIGURE**. Protective effects were measured over 4 days as a function of the length of survival of the cultured cells in the presence of the inventive compounds as compared to survival of the cells in the absence of the inventive compounds. 100% survival (y axis) means that a number of treated cells all survived throughout the test period while an equal number of untreated cells died.

FIGURES 13(a) through (c) show the structure of commercially available isoquinoline structures whose inhibitory effect with respect to production of TNF- α by human monocytes prior to modification to add side chain substituents according to the invention was tested. Except for Compound S52,626-6 (6,7-dimethoxy- 1(2H)-isoquinoline, which possessed mild inhibitory activity as shown in **FIGURE 10**) none of the tested compounds possessed any such inhibitory activity prior to their modification according to the invention, even at concentrations up to 500 μ M.

DETAILED DESCRIPTION OF THE INVENTION

Unless otherwise noted, each of the references cited in the following disclosure is incorporated herein by reference for the purpose of illustrating the level of knowledge and skill in the art with respect to the subject matter discussed. In view
5 of this disclosure, those of ordinary skill in the art will be familiar with, or can readily ascertain, equivalents to the techniques, compounds and other material discussed in each reference suitable for use in the invention.

I.

Composition and Activity 10 of the Bifunctional Activation Cascade Antagonists

The bifunctional activation cascade antagonists comprise a JNK activation cascade antagonist constituent as described in commonly owned US Patent Application Serial Nos. 08/858,778, 08/367,102 and 08/482,551 (which describe the constituent compounds but do not suggest their selectivity for antagonizing the JNK
15 activation cascade) and another anti-inflammatory compound which antagonizes the p38 activation cascade. Each constituent inhibits PIC production, especially production of $\text{TNF}\alpha$. Examples of JNK and p38 activation cascade antagonists are identified in Table I and compared to other anti-inflammatory agents. In the Table, the CSAID compounds which begin with "SK&F" or "SB" are manufactured by
20 Smith-Kline

Beecham Pharmaceuticals, USA. The "FR" compound is manufactured by Fuji Laboratories, Japan. The "SR" compound is manufactured by Sanofi Recherche, France. Other compounds listed are commercially available from different sources.

5 It should be noted that the term "CSAID" is used in this disclosure as a generic term for p38 activation cascade antagonists, especially pyridylimidazole compounds. However, it is acknowledged that CSAID is also used as a trademark by SmithKlineBeecham Pharmaceuticals for specific pyridylimidazole compounds which antagonize p38 activation. Such compounds are included in, but do not exclusively comprise, the class of compounds referred to herein as "CSAIDs".

TABLE I

COMPOUND CLASS	ACTIVITY	EXAMPLES
JNK activation cascade antagonists	PIC inhibitors, affect JNK activation cascade	Described in US Patent Application Serial Nos.08/858,778, 08/367,102 and 08/482,551
CSAIDS	PIC inhibitors, affects p38 activation cascade	SK&F 86002 (also inhibits CO/5-LO), SK&F 104351, SK&F 105561, SK&F 105809, SK&F 202190, SB 203580, SB 206718, PD 098059 SR31747 FR133605
5 NSAID prostanoglandin endoperoxidase (PGHS) and 5-lipoxygenase inhibitors (combination compounds also known as cyclooxygenase(CO)/lipoxygenase (5-LO) dual inhibitors)	inhibit CO and/or 5-LO; tend to stimulate PIC production	ibuprofen naproxen
10 corticosteroids	PIC inhibitor	cortisone, fluticasone propionate, beclomethosone, prednisolone
phosphodiesterase (PDE) inhibitors	inhibit PDE in signalling transduction pathways	rolipram (inhibits 4-PDE) RO-1724 (same target activity)

Of the classes of compounds listed in Table I, the NSAIDs and PDE inhibitors have, respectively, undesirable PIC stimulatory activity and side-effects. Similarly, while useful, corticosteroids pose the risk of certain undesirable side-effects

and are not as specific in their PIC inhibitory activity as CSAIDs. For this reason, pyridylimidazole compounds, especially CSAIDs, are the compounds of choice for use as partners to JNK activation cascade antagonists to form the bifunctional activation cascade antagonists of the invention, which may be administered alone or
5 together with other anti-inflammatory agents and medicaments as indicated.

Administered alone, CSAIDs inhibit 70-80% or less of $\text{TNF}\alpha$ release in an art-accepted model of stress-induced inflammation (Example 2). CSAIDs predominantly antagonize activation of p38 kinase, but do not affect JNK activation to a significant degree. Conversely, the activity of the JNK activation cascade antagonist
10 constituents of the bifunctional activation cascade antagonists of the invention is predominantly directed toward inhibition of PIC-stimulated JNK activation. These constituents of the inventive compounds have little impact on p38 activation.

Surprisingly, combination of CSAIDs with JNK activation cascade antagonists produces a synergistic effect which gives the resulting bifunctional
15 activation cascade antagonists greater anti-inflammatory effect than is achieved by either compound alone. For example, as measured by suppression of $\text{TNF}\alpha$ release in stressed cells, up to 100% inhibition is achieved by bifunctional activation cascade antagonists (Example 1).

The JNK activation cascade antagonist and CSAID constituents of the
20 bifunctional activation cascade antagonists of the invention antagonize activation of different MAPK pathways in a non-overlapping manner. While the invention is not to be limited by any particular theory as to its mechanism of action, it is likely that the increased potency of bifunctional activation cascade antagonists as compared to their


constituent components is owing to a synergism produced by their non-overlapping effects on both of the PIC-sensitive MAPK signalling pathways. Thus, the bifunctional activation cascade antagonists of the invention exert synergistically enhanced, non-additive, pathway-specific effects on the JNK and p38 activation cascade pathways. As such, the compounds can provide greater anti-inflammatory

5 potency, with relatively low risk of additive toxicity, than would be expected from a combination of anti-inflammatory agents.

Descriptions of representative CSAIDs and JNK activation cascade antagonists useful as constituents of the bifunctional activation cascade antagonists of the invention are provided in Section II, A and B of this disclosure. For example,



10 CSAID SB 203580 (inhibits inflammation in several animal models; Badger, *et al.*, *J.Pharmacol.Exp.Ther.*, 3:1453-1461, 1996) and SB 210313 (Boehm, *et al.*, *J.Med.Chem.*, 39:3929-3927, 1996). Particularly potent JNK activation cascade antagonist constituents are compounds 54 and 54a (morpholinyl esters of Compound

15 52, described in Example XVII; isoquinilones), compound 6 (a purine) or compound 37 (a pteridine). Representative structures for each of these high potency constituents are:




CSAID SB203580

(2)



COMPOUNDS 54 and 54a


(3)




Where R_2 is $(CH_2)_3 COOEt$

COMPOUND 6

(4)



CH_3CHO



5

COMPOUND 37

The constituents of the bifunctional activation cascade antagonists may be co-administered as separate compounds. However, to ameliorate their systemic activity, the constituents may be conjugated to one another by a bond which is severable *in vivo* and delivered as a prodrug or combined as separate constituents within a single unit delivery vehicle, such as a liposome.



10

Suitable conjugation means include amide, ester and also linkages, which are separable *in vivo* by, for example, the action of proteinases or esterases. A particularly useful conjugation bond that is cleavable *in vivo* is a tertiary N acyloxymethyl amide bond (Moreira, *et al.*, *Tetrahedron Lett.*, 35:7107-7110 (1994),
5 the disclosure of which is incorporated herein to illustrate the steps for formation of such bonds).


Briefly, tertiary N acyloxymethyl amide bonds are formed to stabilize carboxylic acid or secondary amide moieties on prodrugs (i.e., drugs bound to inert carriers by a bond which is separable *in vivo*). The bonds involve direct coupling of a
10 $R_1CoNR_2CH_2$ (secondary amide) moiety to an R_3CO_2H drug.

Examples of bifunctional activation cascade antagonists bound by ester-linkages are:


(5)



SB203313/COMPOUND 54 or 54a



SB203580/COMPOUND 37



SB203313/COMPOUND 37

5 wherein, for all compounds, n is any number of carbon atoms from 1 to 7,
 O or N;

 R₁, if present, is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or
 aralkyl having less than 7 carbon atoms;

 R₆ and R₇ are H, OH or OR₁, in any combination;

10 Z is N or C;

 X, where Z is C, is H, halogen, N₃, NO, NH₂, NHR₁, N(R₁)₂ or COR₁; and,

 A is H, halogen, N₃, NO, NH₂, NHR₁, N(R₁)₂ or COR₁.

The bifunctional activation cascade antagonists of the invention may be prepared in a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers preferred for use with the bifunctional activation cascade antagonists of the invention may include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/ aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, antioxidants, chelating agents, and inert gases and the like. A composition of bifunctional activation cascade antagonists may also be lyophilized using means well known in the art, for subsequent reconstitution and use according to the invention.

Absorption promoters, detergents and chemical irritants (e.g., keratinolytic agents) can enhance transmission of a bifunctional activation cascade antagonist to a target tissue. For reference concerning general principles regarding absorption promoters and detergents which have been used with success in mucosal delivery of organic and peptide-based drugs, see Chien, *Novel Drug Delivery Systems*, Ch. 4 (Marcel Dekker, 1992).

Examples of suitable nasal absorption promoters in particular are set forth at Chien, *supra* at Ch. 5, Tables 2 and 3; milder agents are preferred. Suitable agents

for use in the method of this invention for mucosal/nasal delivery are also described in Chang, *et al.*, *Nasal Drug Delivery*, "Treatise on Controlled Drug Delivery", Ch. 9 and Table 3-4B thereof, (Marcel Dekker, 1992). Suitable agents which are known to enhance absorption of drugs through skin are described in Sloan, Use of Solubility
5 Parameters from Regular Solution Theory to Describe Partitioning-Driven Processes, Ch. 5, "Prodrugs: Topical and Ocular Drug Delivery" (Marcel Dekker, 1992), and at places elsewhere in the text. All of these references are incorporated herein for the sole purpose of illustrating the level of knowledge and skill in the art concerning drug delivery techniques.

10 A colloidal dispersion system may be used for targeted delivery of the bifunctional activation cascade antagonist to an inflamed tissue. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A particularly convenient and effective colloidal system is a liposome.

15 Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. Compounds can be encapsulated within the aqueous interior and be delivered to cells in an active form. In addition to
20 mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells.

 The composition of a liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination

with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations. Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, 5 phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

10 The targeting of liposomes can be classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-
15 endothelial system (RES) in organs which contain sinusoidal capillaries. Active targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand such as a monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell types other than the naturally occurring sites of
20 localization.

 The surface of the targeted delivery system may be modified in a variety of ways. In the case of a liposomal targeted delivery system, lipid groups can be

incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various well known linking groups can be used for joining the lipid chains to the targeting ligand (see, e.g., Yanagawa, *et al.*, *Nuc.Acids Symp.Ser.*, 19:189 (1988); Grabarek, *et al.*, *Anal.Biochem.*, 185:131 (1990); Staros, *et al.*, *Anal.Biochem.*, 156:220 (1986) and Boujrad, *et al.*, *Proc.Natl.Acad.Sci.USA*, 90:5728 (1993)). Examples of targeting ligands are receptor ligands, antigens, enzymes and monoclonal antibodies. Such targeting ligands can be prepared according to conventional techniques (e.g., peptide synthesis) and many are commercially available.

II.

Bifunctional Activation Cascade Antagonist Constituents*JNK Activation Cascade Antagonist Constituents*





JNK activation cascade antagonist constituents useful in the invention





5 include all of the constituents described in commonly owned US Patent Application Serial Nos. 08/858,778, 08/367,102 and 08/482,551, whose disclosures are incorporated herein. For ease of reference, details of the structures of each of these compounds, as well as additional compounds of similar activity, as well as methods for their synthesis, are provided below.





10 JNK activation cascade antagonists of the invention generally comprise purines, pteridines, thiadiazolopyrimidines and quinazolines prepared according to the schemes described below. For reference, the techniques used in synthesizing the compounds are adaptations of the well-known Traube Synthesis protocol (Lister, "Purines" (Wiley-Interscience, 1971), at p. 220), beginning with 4,5-
15 diaminopyrimidines; to wit: (1) for the purines in general, see Brown, "The Chemistry of Heterocyclic Compounds: Fused Pyrimidines", Part II, The Purines, 1971), at pp. 31-90; (2) for the 9-dieazapurines in particular, see Fox, et al., J. Org. Chem., 43:2536, 1978; (3) for the pteridines, see, for a description of the standard Timmis reaction, Nishigaki, et al., Heterocycles, 15:757-759, 1981; Timmis, Nature, 164:13
20 9, 1949, (or other standard Traube-like protocols for preparing pteridines by ring closure of diaminopyrimidines using a two carbon reagent); and, (4) for the pyrimidines, see, Schrage and Hitchings, J. Org. Chem., 16:207, 1951.





JNK activation cascade antagonists, intermediates and compounds tested for comparison of activity to the compounds of the invention are identified in the discussion below by the numbers assigned to each compound in Table 2.





TABLE 2





Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp(°C)	formula	analysis	TNFAIC ₅₀
1	I		102-103			85
2	I		208-210			>200
4	I		162-165	C ₁₁ H ₁₄ N ₄ O ₄	C, H, N	200
5	I		76-78	C ₁₂ H ₁₆ N ₄ O ₄	HRMS	10






Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp (°C)	formula	analysis ^a	TNF α IC ₅₀ ^b
6	I		73-75	C ₁₃ H ₁₈ N ₄ O ₄	C, H, N	6
7	I		84-85	C ₁₄ H ₂₀ N ₄ O ₄	C, H, N	11
8	I		158-160	C ₁₆ H ₁₆ N ₄ O ₄	C, H, N	>200
10	I		101-102	C ₁₃ H ₁₇ BrN ₄ O ₄	C, H, N	60

Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp(°C)	formula	analyses ^a	TNFAIC ₅₀ ^b
11	I		218-219	C ₁₆ H ₁₅ BrN ₄ O ₄	C, H, N	>200
12	I		202-206	C ₁₃ H ₁₈ N ₄ O ₄ S	C, H, N	100
13	I		>293 dec	C ₁₇ H ₁₈ N ₄ O ₄ S	C, H, N	>200
20	V		>218 dec	C ₁₂ H ₂₀ N ₄ O ₃	C, H, N	ND ^c





Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp (°C)	formula	analysis ^a	TNFAIC ₅₀ ^b
24	I		>295 dec	C ₁₂ H ₁₈ N ₄ O ₃	C, H, N	50
25	I		>320			>200
31	I		oil	C ₁₅ H ₂₂ N ₄ O ₄	C, H, N	5
33	V		>270 dec	C ₅ H ₈ N ₄ O ₂	C, H, N	ND





Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type	Structure	mp ($^{\circ}$ C)	formula	analysis ^a	TNFAIC ₅₀ ^b
34	II		287-289	C ₇ H ₆ N ₄ O ₂	C, H, N	ND
35	II		187-189	C ₉ H ₁₀ N ₄ O ₄	C, H, N	ND
36	II		90-92	C ₁₃ H ₁₆ N ₄ O ₄	C, H, N	12
36a	II		199-202	C ₁₁ H ₁₂ N ₄ O ₄	C, H, N	>200

Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp (°C)	formula	analysis ^a	TNF α IC ₅₀ ^b
37	II		53-55	C ₁₅ H ₂₀ N ₄ O ₄	C, H, N	5
37a	II		97-99	C ₁₃ H ₁₆ N ₄ O ₄	C, H, N	>200
38	II		118-119	C ₁₂ H ₁₂ N ₄ O ₄	HRMS	25
39	II		92-95	C ₁₇ H ₂₂ N ₄ O ₆	C, H, N	ND




Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp (°C)	formula	analyses ^a	TNPaIC ₅₀ ^b
40	II		218-222	C ₁₁ H ₁₄ N ₄ O ₂	C, H, N	ND
41	II		66-68	C ₁₇ H ₂₄ N ₄ O ₄	C, H, N	130
41a	II		161-162	C ₁₅ H ₂₀ N ₄ O ₄	C, H, N	ND
42	II		>307 dec	C ₁₃ H ₁₀ N ₄ O ₄	N/A	ND

Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp(°C)	Formula	analyses ^c	TNFAIC ₅₀ ^b
43a	II	 <chem>CCCCN1C(=O)N(C(=O)OCCCC)C2=NC=CC=N12</chem>	N/A	N/A	N/A	N/A
43b	II	 <chem>CCCCCN1C(=O)N(C(=O)OCCCC)C2=NC=CC=N12</chem>	N/A	N/A	N/A	N/A
43c	II	 <chem>CC(C)CN1C(=O)N(C(=O)OCCCC)C2=NC=CC=N12</chem>	N/A	N/A	N/A	N/A
43d	II	 <chem>C1CCOCC1N2C(=O)N(C(=O)OCCCC)C3=NC=CC=N23</chem>	N/A	N/A	N/A	N/A

Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp ($^{\circ}$ C)	formula	analyses ^a	TNFAIC ₅₀ ^b
43e	II		N/A	N/A	HRMS	N/A
43	II		99-100	C ₁₉ H ₂₀ N ₄ O ₄	HRMS	>200
44	III		73-75	C ₁₂ H ₁₈ N ₄ O ₂ S	C, H, N	75
45	III		210-213	C ₅ H ₄ N ₄ O ₂ S	C, H, N	ND

Physicochemical Data for all New Compounds and TNF- α Inhibition Data						
Comp	Type		mp (°C)	formula	analysis ^a	TNPaIC ₅₀ ^b
46	III		142-144	C ₇ H ₈ N ₄ O ₂ S	C, H, N	ND
47	III		45-47	C ₁₁ H ₁₄ N ₄ O ₄ S	C, H, N	25
47a	III		121-122	C ₉ H ₁₀ N ₄ O ₄ S	HRMS	>200
48	III		o.l.l	C ₁₃ H ₁₈ N ₂ O ₃	C, H, N	18


Physicochemical Data for all New Compounds and TNF- α Inhibition Data

Comp	Type		mp(°C)	formula	analyses ^a	TNFAIC ₅₀ ^b
51	IV		oil	C ₁₄ H ₂₀ N ₄ O ₃	C, H, N ^f	ND
52	IV		oil	C ₁₅ H ₁₈ N ₂ O ₄	C, H, N ^g	55
52a	IV		163-165	C ₁₃ H ₁₄ N ₂ O ₄	C, H, N	>200

^aAll compounds analyzed for C, H, N or by exact mass high resolution mass spectrometry; results were within $\pm 0.4\%$ of theoretical values. ^bConcentration of compound in μM which inhibited the product of TNF α by 50% of control. ^cND = not determined. ^dC: calc'd, 58.61; found 59.22. ^eH: calc'd, 4.13; found 3.70. ^fN: calc'd, 10.25; found 9.77. ^gC: calc'd, 59.62; found, 60.61. Compound types are: I=purines; II=pteridines; III=thiadiazolopyrimidines; IV=quinazolines; and, V=isoquinolones. "N/A" means data not available (but can be obtained through conventional analysis techniques).

Purines

Purines JNK activation cascade antagonists have the general formula (8):



where Z is N or CH and R₁ is (CH₂)_nA, where:

A is NH₂, acyloxy, SO₃H, PO₄H₂, NNO(OH), SO₂NH₂, PO(OH)NH₂,

5 SO₂R, or COOR where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl or benzyl;

n is any number of atoms from 1 to 7 having saturated and/or unsaturated carbon to carbon bonds, which atoms may include an oxygen or nitrogen atom in place of a carbon atom to form, respectively, ether or amino linkages;

10

and, preferably, R₁ is a ω-carboxyalkyl, ω-carboxyalkenyl, or ω-carboxyaryl having from 1 to 8 carbon atoms, wherein the aromatic group further has as a substituent A (as defined above);

R_2 is H, an alkyl (including aliphatic and alicyclic, and heteroalicyclic forms), alkenyl, aralkyl having 1 to 7 carbon atoms or a ω -hydroxyalkyl having from 1 to 7 carbon atoms;

R_3 is the same as R_2 ; and

- 5 X is H, any halogen, OH, SH, OR', or SR', where R' is an alkyl, alkenyl, phenyl or benzyl having from 1 to 4 carbon atoms.


These compounds are synthesized per the below synthesis scheme (as described in further detail in the Examples).

SCHEME 1

(PURINES)


Three basic synthesis protocols were utilized in Scheme I; to wit:

Method A



5 Generally, theobromine was used as the starting material under conditions to ensure N-1 alkylation took place in lieu of O-6 alkylation. Compounds 4 through


8, 10 and 11 (Table I) were prepared by this method. Compounds 10 and 11 in particular were prepared by a variation of the alkylation method in which theobromine was first brominated to give 8-bromotheobromine (compound 9), then alkylated. The 8-bromo substituent was also displaced by NaSH to yield the corresponding 8-thioxo derivatives, compounds 12 and 13.



This method is essentially based on the Traube purine synthesis protocol referred to *supra*. The method was used to prepare 1,3,8-trisubstituted xanthines bearing no alkyl group at the N-7 position. In this procedure, the N-1 substituted pyrimidine was alkylated at position N-3. Formation of the purine ring was complete

by nitrosation, reduction of the nitroso to the amine by catalytic hydrogenation, then ring closure using urea or potassium ethyl xanthate to provide compounds 24 and 25 (respectively, 8-oxo and 8-thioxo derivatives). A detailed description of this protocol is provided in the Examples.

5. Method C




The above method was utilized to prepare the N-3 propylpurines. The starting material used was n-propyl urea condensed with ethyl cyanoacetate in the presence of sodium ethoxide to yield the 6-amino-1-propylpyrimidinedione in moderate yield. Commercially available 3-n-propylxanthine could also be used as the starting material.

Ring closure was accomplished as described in method A, except that diethoxymethyl acetate was used as the source of carbon in the ring closure step. Sequential alkylations were then performed using alkyl halides to yield the final compound 31 (ethyl 4-(2,3,6,7-tetrahydro-2,6-dioxo-7-methyl-3-n-propyl-1H-purin-1-yl)butanoic acid). A detailed description of this protocol is provided in the Examples.

Pteridines

Pteridine JNK activation cascade antagonists have the general formula (9):



R_1 is $(CH_2)_nA$, where:

A is NH_2 , acyloxy, SO_3H , PO_4H_2 , $NNO(OH)$, SO_2NH_2 , $PO(OH)NH_2$, SO_2R , or $COOR$ where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl or benzyl;

5 n is any number of atoms from 1 to 7 having saturated and/or unsaturated carbon to carbon bonds, which atoms may include an oxygen or nitrogen atom in place of a carbon atom to form, respectively, ether or amino linkages;

10 and, preferably, R_1 is a ω -carboxyalkyl, ω -carboxyalkenyl, or ω -carboxyaryl having from 1 to 8 carbon atoms, wherein the aromatic group further has as a substituent A (as defined above);

R_2 is H, an alkyl (including aliphatic and alicyclic, and heteroalicyclic forms), alkenyl, aralkyl having 1 to 7 carbon atoms or a ω -hydroxyalkyl having from 1 to 7 carbon atoms;

15 R_4 is the same as R_2 , OH or an O-alkyl having from 1 to 5 carbon atoms;

R_5 is the same as R_2 , OH or an O-alkyl having from 1 to 5 carbon atoms;


and,

Z is N or CH.

These compounds are synthesized per the below synthesis scheme (which is described in further detail in the Examples.)


SCHEME II

(PTERIDINES)



- (a) $\text{Na}_2\text{S}_2\text{O}_4, \text{H}_2\text{O}$
 (b) 3,4-hexandione
 (c) $\text{R}_1\text{X}, \text{DMF}, \text{K}_2\text{CO}_3$
 (d) glyoxal:sodium bisulfite
 (e) phenethylamine
 (f) dimethylacetylene dicarboxylate

Method C (above) was chosen as a convenient method to produce N-alkyls in preference to O-alkyls in this group. Method C was modified to this end as follows:



Synthesis of the pteridines was based on orthodiaminopyrimidines as precursors. Ring closure of the orthodiamines (compounds 33 and 28) was accomplished with a two carbon source (e.g., glyoxal) to produce compounds 34 and 35 (N-1 substituted pteridines). Alkylation at N-3 as described with respect to


5 Method A produced the desired pteridines (compounds 36-38). Further, use of 3,4-hexanedione in the ring closure step produced a more lipophilic derivative (compound 41; 6,7-diethyl pteridine). Condensation of compound 22 with dimethylacetylene dicarboxylate formed compound 39 (1,3-dialkylpteridine), while treatment of

10 compound 27 phenethyl amine followed by alkylation provided compound 43 (6-phenyl dialkyl pteridine). Both of the latter protocols utilized a Timmis reaction to produce the desired products. A detailed description of these protocols is provided in the Examples.

Thiadiazolopyrimidines

Thiadiazolopyrimidine JNK activation cascade antagonists have the

15 general formula (10):



R_1 is $(CH_2)_nA$, where:

A is NH_2 , acyloxy, SO_3H , PO_4H_2 , $NNO(OH)$, SO_2NH_2 , $PO(OH)NH_2$, SO_2R , or $COOR$ where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl or benzyl;

5 n is any number of atoms from 1 to 7 having saturated and/or unsaturated carbon to carbon bonds, which atoms may include an oxygen or nitrogen atom in place of a carbon atom to form, respectively, ether or amino linkages; and

and, preferably, R_1 is a ω -carboxyalkyl, ω -carboxyalkenyl, or ω -
10 carboxyaryl having from 1 to 8 carbon atoms, wherein the aromatic group further has as a substituent A (as defined above); and


R_2 is H, an alkyl (including aliphatic and alicyclic, and heteroalicyclic forms), alkenyl, aralkyl having 1 to 7 carbon atoms or a ω -hydroxyalkyl having from 1 to 7 carbon atoms.

15 These compounds are synthesized per the below synthesis scheme (as described in further detail in the Examples).

SCHEME III

(THIADIAZOLOPYRIMIDINES)

Method C (above) was chosen as a convenient method to produce N-alkyls in preference to O-alkyls in this group. Method C was modified to this end as follows:




Synthesis of the pyrimidines was based on orthodiaminopyrimidines as precursors. Ring closure of the orthodiamines was accomplished by treatment with thionyl chloride in the presence of pyridine. Alkylation of these intermediates produced compounds 47 and 48 (disubstituted pyrimidines). A detailed description of this protocol is provided in the Examples.

Isoquinolones

Isoquinoline JNK activation cascade antagonists have the general formula

(11):



5 R_2 is $(CH_2)_nA$, where:

A is NH_2 , acyloxy, SO_3H , PO_4H_2 , $NNO(OH)$, SO_2NH_2 , $PO(OH)NH_2$,
 SO_2R , or $COOR$ where R is H, an alkyl having from 1 to 4 carbon atoms,
 an alkenyl having from 1 to 4 carbon atoms, tetrazolyl or benzyl;

10 n is any number of atoms from 1 to 7 having saturated and/or unsaturated
 carbon to carbon bonds, which atoms may include an oxygen or nitrogen
 atom in place of a carbon atom to form, respectively, ether or amino
 linkages;

and, preferably, R_2 is a ω -carboxyalkyl, ω -carboxyalkenyl, or ω -carboxyaryl having from 1 to 8 carbon atoms, wherein the aromatic group further has as a substituent A (as defined above); and

5 R_3 is H, an alkyl (including aliphatic and alicyclic, and heteroalicyclic forms), alkenyl, aralkyl having 1 to 7 carbon atoms or a ω -hydroxyalkyl having from 1 to 7 carbon atoms;

R_4 is H, OH, NH_2 or O-alkyl having from 1-7 carbon atoms;

R_5 is H, OH, NO, NO_2 , NH_2 an O-alkyl having from 1-4 carbon atoms, or X where;

10 X is H, any halogen, OH, SH, OR' , or SR' , where R' is an alkyl, alkenyl, phenyl or benzyl having from 1 to 4 carbon atoms; and,

R_7 is H, OH, NO, NO_2 , NH_2 an O-alkyl having from 1-7 carbon atoms, or X where:

15 X is H, any halogen, OH, SH, OR' , or SR' , where R' is an alkyl, alkenyl, phenyl or benzyl having from 1 to 7 carbon atoms.

These compounds were synthesized by purchasing isoquinolines from Aldrich Chemical and adding side chains to the ring structure with respect to the compounds of the invention. Only the 6,7-dimethoxy-1(2H)-isoquinoline compound

(Aldrich # S52,626-6) had any inhibitory effect on TNF- α production prior to addition of the side chains described above.

Quinazolines


Quinazoline JNK activation cascade antagonists are synthesized according to the following scheme and are represented in structure by compound 52:

Scheme IV

5

(DIDEAZAPTERIDINES OR QUINAZOLINES)


Compound 52 (a quinazoline derivative; ethyl 4-(1-methyl-2,4-dioxoquinazol-3-yl)butanoic acid) was produced as follows:



The starting material for this protocol was N-methyl isatoic anhydride. A detailed description of this protocol is provided in the examples.

Increasing water solubility

The carboxylic ester and acid derivatives of the compounds described above can be converted to substituted carboxylic esters, where the substituent of the ester increases the water solubility of the compounds.




- 5 Particular substituents that increase the water solubility of the compounds effectively are aminoalkyl esters. The amino group of the aminoalkyl ester preferably is a secondary or tertiary amino group. The amino substituents have between 1 and 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom. The preferred aminoalkyl is a tertiary amine and the amino substituents are preferably are
- 10 ether groups. The most preferred aminoalkyl is N-morpholinoethyl. The carboxylic ester and carboxylic acid derivatives can be converted to the morpholinoethyl ester form by the method shown in Scheme V.

Compound 53 (an isoquinolone derivative; ethyl (6,7-dimethoxy-1(2*H*)-isoquinolonyl)butanoic acid was converted to compound 54, a morpholinoethyl ester (isolated as a hydrochloride salt) as follows:

Scheme V

5

(MORPHOLINOETHYL DERIVATIVES: COMPOUND 54)



CSAID Constituents


CSAIDs are available from, for example, the sources listed in the legend to Table 1. Chemical descriptions and data demonstrating activity for many of these compounds have been published in the art in, for example, US Patent No. 5,658,903; US Patent No. 5,593,992; US Patent No. 5,317,019; US Patent No. 4,794,114; US Patent No. 4,778,806; Published PCT Application No. WO9725045; Published PCT Application WO9725047; Published PCT Application WO9725048; Published PCT Application No. WO9640143; European Patent No. 565582; Austrian Patent No. 586907; Griswold, *et al.*, *Drugs Exp.Clin.Res.*, 19:243-248 (1993)(SK&F 86002, 105809 and 104351; *in vivo*=32, 48 and 34 mg/kg, respectively); Marshall, *et al.*, *Biochem.Pharmacol.*, 42:813-824 (1991) (SK&F 105809; *in vitro* data for inhibition of CO, 5-LO, IL-1 and PGHS); Griswold, *et al.*, *J.Immunol.Methods*, 195:1-5 (1996) (SB 203580); and Badger, *et al.*, *J.Pharmacol.Exp.Ther.*, 279:1453-1461 (1996) (*in vivo* data for SB 203580); Boehm, *et al.*, *J.Med.Chem.*, 39:3929-3937 (1997) (SB 210313)). Each of these references are incorporated herein for ease of reference in identifying CSAIDs for use in the invention.

For further ease of reference, the chemical names of several known CSAIDs are set forth below:






SB 210313: 1-[3-(4-morpholinyl)propyl]-4-(4-fluorophenyl)-5-(4-pyridyl)imidazole


- SK&F 105809: 2-(4-methylsulfinylphenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole
- SK&F 105561: 2-(4-methylthiophenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole
- 5 SB 203580: 4-(4-fluorophenyl)-2-(4-methylsulfinyl)-5-(4-pyridyl)imidazole
- SK&F 86002: 5-(4-pyridyl)-6(4-fluorophenyl)-2,3-dihydroimidazole(2,1-b)thiazole

A representative CSAID synthesis scheme (shown for SB 210313) is:




4-pyridylcarbinol





T.F. Gallagher, et al. *Bioorg. Med. Chem. Lett.*
5, 1171-1176, 1995 and US Patent 5656644.



Although not preferred as partners for JNK activation cascade antagonists in the bifunctional activation cascade antagonists of the invention, corticosteroids and other medicaments, such as antibiotics or different anti-inflammatory agents, may also be administered with the inventive compounds if clinically indicated.

5

III.

Kits for Use in Practicing the Methods of the Invention

For use in the methods described below, kits are also provided by the invention. Such kits may include any or all of the following: a bifunctional activation cascade antagonist; a pharmaceutically acceptable carrier (may be pre-mixed with the bifunctional activation cascade antagonist) or suspension base for reconstituting lyophilized bifunctional activation cascade antagonists; additional medicaments; a sterile vial for each bifunctional activation cascade antagonist and additional medicament (if any), or a single vial for mixtures thereof; device(s) for use in delivering the composition to a host; assay reagents for detecting indicia that the anti-inflammatory sought have been achieved in treated animals and a suitable assay device.

10

15

IV.

Methods for Use of the Inventive Compounds.

The bifunctional activation cascade antagonists of the invention may be administered to a mammalian host to retard cellular responses associated with TNF- α and IL-1 production, release and MPAK system activation. As exemplified herein, the methods of the invention are expected to be of particular use in providing

20

protection against inflammation and associated excess formation of fibrotic tissue. In particular, with reference to demonstrated efficacy for the JNK activation cascade antagonist and CSAID constituents of bifunctional activation cascade antagonists, these compounds can be expected to have therapeutic efficacy in the treatment of

5 arthritis, spondylitis, bone resorption (osteoarthritis), sepsis, septic shock, endotoxic shock, inflammatory bowel disease, asthma, bronchitis, chronic pulmonary inflammatory disease, silicosis, reperfusion injury, graft versus host reactions, allograft rejection, fever, viral and bacterial infection, inflammation of the joints, psoriasis and eczema, radiation dermatitis, cell senescence and apoptosis, multiple

10 sclerosis and other inflammatory conditions.

The bifunctional activation cascade antagonists of the invention are administered to a host using any available method and route suitable for drug delivery, including *ex vivo* methods, as well as systemic or localized routes. Those of ordinary skill in the clinical arts will be familiar with, or can readily ascertain, means for drug

15 delivery, which are briefly discussed below.

The entrance point for many inflammatory stimulants into a host is through the skin or mucosa. Thus, delivery methods and routes which target the skin (e.g., for cutaneous and subcutaneous conditions) or mucosa (e.g., for respiratory, ocular, lingual or genital conditions) will be especially useful.

20 Intranasal administration means are particularly useful in addressing respiratory inflammation, particularly inflammation mediated by antigens transmitted

from the nasal passages into the trachea or broncheoli. Such means include inhalation of aerosol suspensions or insufflation. Nebulizer devices suitable for delivery of drug compositions to the nasal mucosa, trachea and bronchioli are well-known in the art and will therefore not be described in detail here. For general review in regard to
5 intranasal drug delivery, those of ordinary skill in the art may wish to consult Chien, *Novel Drug Delivery Systems*, Ch. 5 (Marcel Dekker, 1992).

Dermal routes of administration, as well as subcutaneous injections, are useful in addressing allergic reactions and inflammation in the skin. Examples of means for delivering drugs to the skin are topical application of a suitable
10 pharmaceutical preparation, transdermal transmission, injection and epidermal administration.

For transdermal transmission, absorption promoters or iontophoresis are suitable methods. For review regarding such methods, those of ordinary skill in the art may wish to consult Chien, *supra* at Ch. 7. Iontophoretic transmission may be
15 accomplished using commercially available "patches" which deliver their product continuously via electric pulses through unbroken skin for periods of several days or more. Use of this method allows for controlled transmission of pharmaceutical compositions in relatively great concentrations, permits infusion of combination drugs and allows for contemporaneous use of an absorption promoter.

20 An exemplary patch product for use in this method is the LECTRO PATCH trademarked product of General Medical Company of Los Angeles, CA. This product electronically maintains reservoir electrodes at neutral pH and can be adapted to provide dosages of differing concentrations, to dose continuously and/or to dose periodically. Preparation and use of the patch should be performed according to

the manufacturer's printed instructions which accompany the LECTRO PATCH product; those instructions are incorporated herein by this reference.

Opthalmic administration (e.g., for treatment of allergic conjunctivitis) involves invasive or topical application of a pharmaceutical preparation to the eye.

- 5 Eye drops, topical cremes and injectable liquids are all examples of suitable mileaus for delivering drugs to the eye.

- Systemic administration involves invasive or systemically absorbed topical administration of pharamaceutical preparations. Topical applications as well as intravenous and intramuscular injections are examples of common means for systemic
10 administration of drugs.

- The compounds of the invention vary in potency. A summary of the potency of each JNK activation cascade antagonist (expressed as a percentage of inhibition of intracellular responses to LPS, namely, the production of TNF- α , where responses to pure LPS=100% and are measured as the concentration of the inventive
15 compound needed to inhibit TNF- α production by 50%) is provided in Table 2, elsewhere above.

- The relative potencies of other antiinflammatory constituents, including CSAIDs, are known in the art. For example, the IC₅₀ (level at which 50% of TNFa release is inhibited) for CSAIDs are published (see, e.g., Griswold, *et al.*, *Drugs*
20 *Exp.Clin.Res.*, 19:243-248 (1993)(SK&F 86002, 105809 and 104351; *in vivo*=32, 48 and 34 mg/kg, respectively); Marshall, *et al.*, *Biochem.Pharmacol.*, 42:813-824 (1991) (SK&F 105809; *in vitro* data for inhibition of CO, 5-LO, IL-1 and PGHS); Griswold, *et al.*, *J.Immunol.Methods*, 195:1-5 (1996) (SB 203580); and Badger, *et al.*,

J.Pharmacol.Exp.Ther., 279:1453-1461 (1996) (*in vivo* data for SB 203580); Boehm, *et al.*, *J.Med.Chem.*, 39:3929-3937 (1997) (SB 210313)).

The maximum potency of any of these compounds administered alone is about 70-80% TNF α inhibition (Example I). Those of ordinary skill in the art will
5 recognize that lesser or greater dosages of the compounds of the invention may be required depending on the potency of the particular compound being administered.

Dosages of the compounds of the invention will vary depending on the age, weight and presenting condition of the host to be treated, as well as the potency of the particular compound administered. Such variables will readily be accounted for
10 by those of ordinary skill in the clinical art. In particular, dosages will be adjusted upward or downward for each recipient based on the severity of the condition to be treated and accessibility of the target cells to the pharmaceutical formulations of the invention. Where possible, it will be preferable to administer the pharmaceutical formulations of the
15 invention locally at the site of the target cells; e.g., into inflamed skin or by infusion to another organ of the host. Thus, dosages will also vary depending on the route of administration and the extent to which the formulations of the invention are expected to reach target cells before dilution or clearance of the formulation.

Taking these factors into account, acceptable CSAID and JNK activation
20 cascade antagonist dosage ranges for each compound administered separately are about 0.1 to 80 mg/kg/day for oral and parenteral routes; about 0.1-150 mg/kg/day for topical routes and about 0.01-1 mg/kg/day for respiratory routes. Due to their

enhanced potency, bifunctional activation cascade antagonists are administered in lower dosages (e.g., at about a 10-40% lower dosage). Those of ordinary skill in the clinical arts will be able to determine medically sound dosing schedules for patients with particular presenting conditions, taking into account the severity of the condition, 5 the patient's overall health, patient age and weight, and other clinically relevant factors. These dosages may be combined with other conventional pharmaceutical therapies for inflammation and fibrosis; e.g., corticosteroids.

Those of ordinary skill in the art will be familiar with, or can readily ascertain, the identity and clinical signs of specific inflammatory conditions, and can 10 identify clinical signs of improvement therein (such as reductions in serum levels of TNF- α and improvement in clinical health) in addition to those exemplified herein.

V.

Methods for Identification of Therapeutically Effective Analoguesof Bifunctional activation cascade antagonists

- Those of ordinary skill in the art will be familiar with means to develop analogues to the bifunctional activation cascade antagonists and constituents specifically described herein which, although not structurally identical thereto, may possess the same biological activity. Such compounds are within the scope of the invention and may be identified according to the protocols described below and in the Examples.
- Though exposure of cells to the compounds of the invention under controlled conditions, the responsiveness of cells to inflammatory agents and intracellular mechanisms therefor can be investigated. This information will not only better elucidate the intracellular pathways responsible for cellular responses to particular stimuli, but will also aid in the identification of anti-inflammatory and anti-fibrosis therapeutic compounds.

- To identify and select therapeutic compounds for use in treating conditions such as inflammation and fibrosis, lymphocytes, monocytes, neutrophils, intracellular components such as microsomes or immunologically naive animals are exposed to a PIC and the candidate therapeutic compound. Specifically, a control is incubated with a known amount of the inflammatory or fibroblast proliferation inducing agent. Treatment groups are exposed to the same amount of inflammatory or fibroblast proliferation inducing agent as well as aliquots of the candidate therapeutic compound. Inflammatory responses or fibroblast proliferation in each group are

detected by conventional means known to those of skill in the art (such as the assay steps described in the examples) and compared.

The invention having been fully described, examples illustrating its practice are set forth below. These examples should not, however, be considered to

5 limit the scope of the invention, which is defined by the appended claims.

In the examples, the abbreviation "min." refers to minutes, "hrs" and "h" refer to hours, and measurement units (such as "ml") are referred to by standard abbreviations. "mp" refers to melting point.

EXAMPLE I

INHIBITION OF PIC ACTIVATION OF THE MAPK SYSTEM
BY BIFUNCTIONAL ACTIVATION CASCADE INHIBITORS

As an *in vitro* model of MAPK system activation, peripheral blood

5 monocyte cells (PBMC) were obtained from healthy human donors and stimulated with a PIC-sensitive-MAPK activator, namely lipopolysaccharide (LPS, Sigma). TNF α production was measured as the hallmark of MAPK cascade activation antagonism.

The PBMC were isolated from heparinized normal human blood by ficoll-
10 histopaque density centrifugation. The cells were plated in 96 well plates with a JNK activation cascade antagonist (compound 54 or compound 37), a CSAID (SB 210313) or a bifunctional activation cascade antagonist composed of compound 54 or 37 and SB 210313. The latter was synthesized as described in Boehm, *et al.*, *J.Med.Chem.*, 39:3929-3937 (1997). SB210313 is 1-[3-(4-morpholinyl)propyl]-4-(4-fluorophenyl)-5-
15 (4-pyridyl) and is crystallized from acetone/hexane and stored as a dry powder at room temperature. The synthesized SB210313 compound tested is referred to below as I-105.

Each compound tested was added to the wells in equal volumes at concentrations of 10 μ m or 50 μ m. The plated cells were suspended in RPMI 1640
20 supplemented with 20% autologous plasma and glutamine and plated at a density of 5X10⁵ cells/ml. Cells were incubated in the plates for 60 minutes then each well received LPS at a final concentration of 10 μ g/ml. 18 hours later, 100 μ l of each well

was collected and assayed for TNF α content using a commercially available mouse anti-human TNF α antibody (Upstate Biotechnology, Inc.).

In the TNF α inhibition assay, compound 54 achieved 50% inhibition (IC₅₀) at 10 μ m. The IC₅₀ for compound 37 was achieved at 20-25 μ m. The IC₅₀ for
5 SB210313, one of the most potent CSAIDs, was achieved at 1.6 to 3.2 μ m. However, the maximal level of inhibition achieved by any of the compounds was only 70-80%.

In startlingly contrast, the bifunctional activation cascade antagonist composed of compound 37 (the JNK activation cascade antagonist of *lesser* potency as compared to compound 54) and SB 210313 achieved IC₅₀ in as little as 0.78 μ m, a
10 level more than merely additive of the activities of each constituent alone. Thus, the bifunctional activation cascade antagonists of the invention achieve synergy between the JNK activation cascade antagonist and CSAID constituents.

EXAMPLE II

EFFECT OF JNK ACTIVATION CASCADE ANTAGONISTS AND CSAIDs 15 ON THE JNK ACTIVATION CASCADE

To demonstrate the probable efficacy of bifunctional activation cascade antagonists in antagonizing activation of JNK, the data below demonstrating the efficacy of JNK activation cascade antagonists in this context are provided.

MOLT-4 human lymphoblastoid cells were used as models of JNK
20 activation in the presence of anisomycin.

MOLT-4 cells were preincubated with compound 54 to demonstrate the effect of compound 54 on JNK activation. The compound was dissolved in phosphate buffered saline and provided to the cells in concentrations of 50 μ m or 10 μ m. Incubations were made for intervals of 15, 30 or 60 minutes. Each incubated cell sample was activated with 50nM anisomycin for 30 minutes. Cells exposed only to PBS served as a control and, for comparison, one group of cell samples were exposed to compound 54 alone (without anisomycin) or anisomycin alone (without compound 54).

After the activation period, the MOLT-4 cell samples were washed three times and lysed. 100 μ g protein samples were used as reagents to measure enzyme activity in the MOLT-4 cell samples by a standard binding kinase assay, using GST c-Jun as the ligand and substrate.

For comparison, RAW mouse macrophages were stressed with a different stressor and the inhibitory potency of compound 54 vis-a-vis JNK activation was tested. The cells were pre-incubated with compound 54 for 60 minutes at increasing concentrations and activated with 1 μ g/ml LPS for 30 minutes. The cells were washed three times and lysed before testing.

The IC₅₀s for compound 54 varied from 10-50 μ m, depending on the stressor, the time of preincubation and the cell line studied. Specifically, the IC₅₀ for compound 54 with respect to anisomycin-induced JNK activation was achieved in MOLT-4 cells at 10 μ m after 60 minutes of incubation. In contrast, virtually no

change in JNK activation is achieved by CSAIDs in similar models (see, e.g., Kumar, *et al.*, *Biochem. Biophys. Res. Commun.*, 235:533-538 (1997)).

As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE III

EFFECT OF CSAIDs AND JNK ACTIVATION CASCADE ANTAGONISTS ON THE p38 ACTIVATION CASCADE

To demonstrate the probable efficacy of bifunctional activation cascade antagonists in antagonizing activation of p38, the data below demonstrating the activity of JNK activation cascade antagonists and of CSAIDs in this context are provided.

Lymphocyte and monocyte extracts were prepared under similar conditions and were immunoprecipitated with affinity purified goat polyclonal antibody against the 20 amino acid N terminus of p38 (Santa Cruz Biotechnology), together with protein A SEPHAROSE™. The immunoprecipitates were washed, mixed with glutathione S-transferase-ATF2 SEPHAROSE™ beads as substrate, and resuspended in kinase buffer containing [γ -³²P]ATP. Samples were processed as described in the preceding Example, separated by SDS-PAGE and analyzed by autoradiography with Coomassie blue staining. Densitometry tracings of the 49 kDa GST-ATF2 band

showed that compound 54 does not block p38 kinase activation to an appreciable extent (only about 20% inhibition was achieved at 100 μ m).

In contrast, similar assays have demonstrated that CSAIDs inhibit activation of p38, but not of other kinases (see, e.g., Wilson, *et al.*, *Chem.Biol.*, 4:423-
5 431 (1997)).

These data, and those set forth in Example II, demonstrate that the JNK activation cascade antagonist and CSAID constituents of the bifunctional activation cascade antagonists of the invention antagonize activation of different MAPK pathways in a non-overlapping manner.

10 As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE IV

INHIBITION OF CELL SENESENCE AFTER SERUM DEPRIVATION
IN SERUM-DEPENDENT CELLS BY JNK ACTIVATION CASCADE
ANTAGONISTS

5 To demonstrate the probable efficacy of bifunctional activation cascade antagonists in inhibiting cell senescence, the data below demonstrating the efficacy of JNK activation cascade antagonists in this context are provided.

 Many cell types are dependent upon serum factors for growth. Thus, deprivation of such cells of serum provides a model for assessment of compounds to
10 modulate cell responses to intracellular ceramide-mediated signal transduction. In particular, withdrawal of serum from serum-dependent cell cultures produces increased intracellular levels of endogenous ceramide and may also increase intracellular levels of endogenous diacyl glycerol (see, e.g., Jayadev, *et al.*, *J.Biol.Chem.*, 270:2047-2052, 1995).

15 To evaluate the inhibitory effect of the compounds of the invention on ceramide-associated conditions *in vitro*, the serum withdrawal model was used. Specifically, 3T3 fibroblast cells were seeded in 96 well microtiter plates in DMEM in the presence of 10% fetal bovine serum. The cells were incubated to 90% confluence.

 The medium was removed, the cells washed and reincubated in serum-free
20 DMEM. Compound no. 37 and cell permeable ceramide were added to the wells at concentrations of, respectively, 0, 4, 40 or 400 μ M compound no. 37 and 0, 5 or 10 μ M of ceramide. After 24 hrs. incubation, 0.5 μ Ci of [3 H] thymidine was added to

each well for 2 hrs. DNA synthesis in the tested cell population was assessed by conventional techniques for detection of [³H] thymidine incorporation. The results of this assay are indicated in FIGURE 1 and establish the cell senescence inhibitory efficacy of the inventive compounds (as represented by compound no. 37).

- 5 As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE V

INHIBITION OF CELL APOPTOSIS AFTER CD95 STIMULATIONBY JNK ACTIVATION CASCADE ANTAGONISTS

To demonstrate the probable efficacy of bifunctional activation cascade
5 antagonists in inhibiting cell apoptosis, the data below demonstrating the efficacy of
JNK activation cascade antagonists in this context are provided.

Engagement of cell surface receptor CD95 (also known as Fas/Apo-1
antigen) triggers cell apoptosis. DX2 is a functional anti-FAS (CD95) antibody which
will, on binding of CD95, activate the Smase catalysis of sphingomyelin hydrolysis
10 and production of ceramide (see, re DX2, Cifone, *et al.*, *J.Exp.Med*, 177:1547-1552,
1993, the disclosure of which is incorporated herein by reference for use in accessing
the DX2 antibody). Thus, binding of CD95 is a model for induction of apoptosis via
the sphingomyelin signal transduction pathway.

To assess the inhibitory effect of the compounds of the invention on
15 ceramide-mediated cell apoptosis, human T lymphoblasts (Jurkat) were suspended at
2x10⁶ cells per ml in RPMI-1640 supplemented with insulin, transferrin, selenium and
glutamine. After incubation for 2 hrs. at room temperature with either compound no.
37, compound no. 6, pentoxifylline or a control compound (Ro-1724), 25 ng/ml of
anti-FAS antibody was added to each suspension. After another 2 hrs., cell apoptosis
20 was measured as a function of the number of cells (counted by hemocytometer) that
excluded the vital dye erythrosin B. The results of the experiment are shown in
FIGURE 2 and establish the apoptosis inhibitory efficacy of the compounds of the
invention (as represented by compounds nos. 6 and 37, particularly the latter).

As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

5

EXAMPLE VI

INHIBITION OF THE ACTIVITY OF CERAMIDE ACTIVATED PROTEIN
KINASE BY JNK ACTIVATION CASCADE ANTAGONISTS

To demonstrate the probable efficacy of bifunctional activation cascade antagonists in inhibiting activation of ceramide activated protein-kinase activation (at
10 a site recognized by MAPK kinases), the data below demonstrating the efficacy of JNK activation cascade antagonists in this context are provided.

Ceramide-activated protein kinase (CaPK) is a 97 kDa protein which is exclusively membrane-bound and is believed to serve a role in the sphingomyelin signal transduction pathway. In particular, CaPK is believed to mediate
15 phosphorylation of a peptide derived from the amino acid sequence surrounding Thr⁶⁶⁹ of the epidermal

growth factor receptor (i.e., amino acids 663-681). This site is also recognized by MAPK kinases. Thus, the effect of the JNK activation cascade antagonists on CaPK

activity in cells is indicative of the effect that the compounds exert on signal transduction in the sphingomyelin pathway.

To that end, Jurkat cells were suspended at 2×10^6 cells per ml in RPMI-1640 medium. After incubation for 2 hrs., either compound 37, 20 μ M of ceramide or
5 25 ng/ml of anti-FAS antibody DX2 were added to each suspension and incubated for 15 mins. After centrifugation and washing, the cells were separately homogenized in a dounce homogenizer.

Ceramide kinase levels in each test sample were assayed as described by Liu, *et al.*, *J.Biol.Chem.*, 269:3047-3052, 1994 (the disclosure of which is
10 incorporated herein for reference and use in assaying ceramide kinase). Briefly, the membrane fraction was isolated from each test sample of treated cell homogenate by ultracentrifugation and run on a 10% PAGE gel. The gel was washed with guanidine-HCL, and renatured in HEPES buffer. Then [32 P]-ATP was added to the gel and left there for 10 mins. Thereafter, the gel was extensively washed with 5% TCA.
15 Autophosphorylated kinase was detected by autoradiography.

The results of this assay are indicated in FIGURE 3, and establish the CaPK inhibitory efficacy of the compounds of the invention (as represented by compound 37).

As demonstrated in Example I, the bifunctional activation cascade
20 antagonists of the invention provide the activity of each antagonistic constituent in a

synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE VII

ABSORBANCE OF UVB RADIATIONBY JNK ACTIVATION CASCADE ANTAGONISTS

To demonstrate the probable efficacy of bifunctional activation cascade
5 antagonists in inhibiting absorbance of UVB radiation, the data below demonstrating
the efficacy of JNK activation cascade antagonists in this context are provided.

Radiation (particularly in the UVB wavelength) is a major cause of skin
damage (including apoptosis) in humans. As indicated elsewhere above, the
sphingomyelin signal transduction pathway is believed to be involved in at least the
10 early stages of development of radiation induced dermatoses (including radiation
dermatitis, sunburn and UVB induced immune suppression from radiation damage to
Langerhans cells in the skin- see, e.g., Haimovitz-Friedman, *et al.*, *J.Exp.Med.*,
180:525-535, 1994 (cellular responses to ionizing radiation); and, Kurimoto and
Streilein, *J.Immunol.*, 145:3072-3078, 1992 (cutaneous immune suppression from
15 UVB exposure)). Thus, a compound which will inhibit cell responses to stimulus of
the sphingomyelin signal transduction pathway by radiation and can be administered
topically at the site of exposure would be of great benefit in retarding the damage
associated with radiation exposure (e.g., through exposure to sunlight or radiation).

To assess the radiation absorbing abilities of the compounds of the
20 invention, the ultraviolet spectra of compounds of the invention (nos. 6 and 37, alone,
in combination and as 8-oxo derivatives) were evaluated and compared to those of a
commercially available sunscreen additive (PABA) and isoquinoline. The spectra
were identified using a KONTRON analytical instrument. As indicated in FIGURE 4,

the compounds of the invention (as represented by compounds nos. 6 and 37) absorbed through most of the UVB region, indicating efficacy in absorbing radiation. Surprisingly, a mixture of compound nos. 6 and 37 proved to absorb throughout the UVB region. Thus, given the somewhat greater absorbance characteristics of
5 compound 37 vis-a-vis compound 6, it can be reasonably expected that mixtures of the two in ratios of 1:1 or greater (favoring compound 37) will have substantial synergistic efficacy in absorbing radiation and retarding its effects on cells.

EXAMPLE VIII

10 INHIBITION OF TNF- α PRODUCTION BY JNK ACTIVATION CASCADE ANTAGONISTS(OTHER THAN COMPOUND 54)

To demonstrate the probable efficacy of bifunctional activation cascade antagonists in inhibiting TNF α production, the data below demonstrating the efficacy of JNK activation cascade antagonists in this context are provided.

As shown in FIGURE 5(a), compounds of the invention having N-1 chain
15 lengths from 2-5 carbons are especially useful in inhibiting TNF- α production *in vitro*, while N-1 chain lengths of about 4 carbons (with a terminal ester) appear to be optimal in this respect (as compared to a control compound; FIGURE 5(b)). Further, the esterified compounds were significantly more effective inhibitors of TNF- α production than their carboxylic counterparts. These data were obtained as follows:

20 Peripheral blood mononuclear cells were isolated from normal human blood on Hypaque-Ficoll density gradients. A portion of the isolated cells were further purified by adherence to gelatin coated flasks.

100 μ l aliquots of monocytes were placed onto 96 well microtiter plates at a density of 5×10^5 cells/ml in RPMI-1640 medium containing 10% fetal bovine serum. After incubation for 24 hrs., various concentrations of the test compounds (FIGURE 10) were added to the plated cells in a volume of 100 μ l and incubated for 1
5 hr. After incubation, 1 μ g/ml of LPS was added to each well.

18 hrs. after exposure of the plated cells to LPS, 100 μ l of medium was collected from each well and assayed (by ELISA) for release of TNF- α , using recombinant human TNF as a standard. The sensitivity of the assay ranged from 10-100 pg/ml.

10 As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE IX

15 IN VIVO AND IN VITRO LEUKOPENIA IN RESPONSE TO LPS AND
INHIBITION OF SAME BY JNK ACTIVATION CASCADE ANTAGONISTS

To demonstrate the probable efficacy of bifunctional activation cascade antagonists in inhibiting *in vivo* leukopenia, the data below demonstrating the efficacy of JNK activation cascade antagonists in this context are provided.

20 As shown in FIGURES 6 through 10 and Table 1, the compounds of the invention effectively reduce cellular response to LPS, a known inducer of TNF- α

production. In the presence of ceramide, the inhibitory activity of the compounds of the invention on LPS induced leukopenia (a phenomenon dependent on $\text{TNF-}\alpha$ induced surface expression of the P-selection class of adhesion molecules) was enhanced (FIGURE 6). However, the inhibitory activity of the compounds of the invention was essentially unaffected by diacylglycerol (FIGURE 7), indicating that the mode of action of the compounds of the invention are not dependent on hydrolysis of phosphatidic acid. These data were obtained as follows:

The leukopenia inhibitory capacity of the test compounds was determined by intraperitoneal administration of 0.5 μg of LPS in saline to ICR female mice (age 6-8 weeks; weight 19-23 g). One hour before receiving the LPS, the mice received the test compound by intraperitoneal injection at a dose of 50 mg/kg (in isotonic saline). Two hours after injection of LPS, 200 μl of blood was collected from each mouse into a heparinized tube and the total count of nucleated cells determined in a hemocytometer.

Calcium independent protein kinase activity was measured, using a 1% triton X-100 extract of Jurkat cells ($5 \times 10^8/\text{ml}$). The reaction mixture consisted of 20 mM Tris HCl pH 7.5, 20 mM MgCl_2 , 20 μM ATP containing 200,000 cpm [$\gamma^{32}\text{P}$] ATP, and 50 μM Myelin Basic Protein. The extract was pre-incubated with A) compound 37 B) compound 37 with or without 10 μM ceramide C) ceramide or D) dihydro ceramide for 15 minutes, followed by addition of substrate and ATP, and incubation at 30°C for 5 minutes. The total count of nucleated cells was measured in

a hemocytometer (FIGURE 8). The same protocol was followed to obtain the results shown in FIGURE 7 (with the addition of diacyl glycerol to some of the test mixtures).

An isoquinoline compound of the invention (compound 52) was also tested
5 *in vitro* for its inhibitory efficacy with respect to LPS induced TNF- α production in human cells.

Human macrophages were cultured in 96 well microtiter plates and incubated with LPS. Aliquots of the stimulated cells were then incubated with, respectively, 0.1, 1, 10, 100 or 1000 μ M of compound 52, compound 37 and a
10 commercially available isoquinoline (6,7-dimethoxy-1(2H)-isoquinoline from Aldrich Chemical; labelled S52-626-6 in FIGURE 9) which, like the compounds of the invention, has an oxygen *ortho* to a ring nitrogen but, *unlike* the compounds of the invention, lacks a side chain

substituent as described above. The inhibitory efficacy of each compound was
15 measured as a function of TNF- α reduction in pg/ml. The results of the experiment are indicated in FIGURE 11 and establish that the compounds of the invention (represented by compound 52 and compound 37) have inhibitory efficacy with respect to reduction in LPS induced TNF- α production by human cells. Other isoquinolines tested did not exert inhibitory activity in the absence of the side chain substituents
20 added according to the invention.

These experiments were repeated in human lymphocytes. The protective abilities of the compounds tested is demonstrated by the data in FIGURES 12 and 13.

As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE X

IN VIVO ANTI-INFLAMMATORY EFFICACY OF CSAIDs

The effects of CSAIDs on inflammation in an *in vivo* model of arthritis have been previously demonstrated. The following references are relevant in this regard and are incorporated herein for ease of reference by those in the art:

Boehm, *et al.*, *J.Med.Chem.*, 39:3929-3937 (1996) (SB 210313)

Badger, *et al.*, *J.Pharmacol.Exp.Ther.*, 279:1453-1461 (1996) (SB 203580)

As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE XI

FIBROBLAST PROLIFERATION IN RESPONSE TO LPS AND INHIBITION OF SAME BY JNK ACTIVATION CASCADE ANTAGONISTS

To demonstrate the probable efficacy of bifunctional activation cascade antagonists in inhibiting fibrosis, the data below demonstrating the efficacy of JNK activation cascade antagonists in this context are provided.

As shown in FIGURE 10, PDGF induced fibroblast proliferation was
5 selectively inhibited by the compounds of the invention. In addition, the compounds were shown not to be cytostatic or cytotoxic, insofar as they did not alter EGF-triggered mitogenesis in the cells tested (FIGURE 11).

These data were obtained as follows:

Mouse fibroblast line 3T3 cells (American Type Culture Collection #CCL
10 92) were seeded into 96 well plates in complete medium and allowed to grow to confluence. The medium was then replaced with medium-free serum and the cells incubated for 24 hrs.

The test compounds were then incubated with the cells for 1 hr before addition of 5 ng/ml human PDGF or EGF was added to each well. After another 24
15 hrs, 1 μ Ci of [3 H]-thymidine was added to each well. 4 hrs later the cells were harvested onto glass fiber filters and the cellular incorporation of [3 H]-thymidine was measured by liquid scintillation counting (FIGURES 10 and 11).

As demonstrated in Example I, the bifunctional activation cascade antagonists of the invention provide the activity of each antagonistic constituent in a
20 synergistic manner. Thus, the bifunctional activation cascade antagonists provide the same activity of each constituent alone at greater potency.

EXAMPLE XII

SYNTHESIS OF COMPOUNDS 2, 4-8 AND 10-13

General Alkylation Procedure for Compounds 4-8, 10, 11 (Method A):

Theobromine or 8-bromotheobromine (2 mmol) was combined with
5 anhydrous K_2CO_3 (2.5 mmol) and dry DMF (15 mL) and the mixture was brought to
75°C. The appropriate alkyl halide (2.5 mmol) was added and the mixture was stirred
at 75°C for 2-18 h. The reaction mixture was cooled, poured into water (125 mL) and
extracted with ethyl acetate (2 x 75 mL). The organic layer was dried over
magnesium sulfate and evaporated to yield a colorless oil or white solid which was
10 triturated with ethyl ether. The resulting solid, often analytically pure, may be
purified further if desired by crystallization from a small amount of ethanol. Yields
58-89%. Compounds 15-17, 31, 36-38, 41, 43, 47, and 48 (described below) were
prepared by this same procedure only using the appropriate precursors in place of
theobromine.

15 General Thiation Procedure for Compounds 12 and 13:

The 8-bromoxanthine 10 or 11 (0.25 mmol) was suspended in anhydrous
ethanol (10 mL) and heated to reflux. $NaSH \cdot (H_2O)_x$ (2.5 mmol) was added and the
mixture became clear, green almost immediately. The mixture was stirred under
reflux for 30 min, cooled and evaporated onto silica gel. Flash column
20 chromatography using 5-7% MeOH in CH_2Cl_2 provided a 63% and 75% yield of 12
and 13, respectively as white solids. Note: Compound 13 was found by 1H NMR to
be the ethyl ester due to transesterification under the reaction conditions.

¹H NMR spectra and elemental analyses or exact mass data were consistent with the assigned structures.

EXAMPLE XIII

SYNTHESIS OF COMPOUNDS 24, 25, 31 AND INTERMEDIATES

5 General Procedure for C-Nitrosation of Pyrimidines (Compounds 18-20, 27, and 32):

The pyrimidine (15 mmol) was suspended in N HCl (30 mL) and an aqueous solution of sodium nitrite (20 mmol in 10 mL) was dripped in with stirring over 10 min. The suspension went from off-white to purple almost immediately.

10 Stirring was continued for 1h, pH adjusted to 5 with ammonia water and the purple solid product collected to provide 75-90% yield after drying. The characteristic lack of the C-5 proton in the ¹H NMR was evident for each compound (Table 2).

General Procedure for the Reduction of 5-Nitroso to 5-Amino Pyrimidines (Compounds 21-23, 28, and 33):

15 The 5-nitrosopyrimidine (15 mmol) was suspended in water (50 mL) and heated to 80-90°C. With stirring, sodium hydrosulfite (45 mmol) was added in portions over 5 min. The color quickly changed from purple to light green and stirring was continued an additional 10 min. The mixture was cooled in ice and filtered. The filtered solid was washed with cold water, EtOH and Et₂O to provide the

20 orthodiamine in 70-88% yield as a tan to pale green solid.

Synthesis of 1-n-Hexyl-3-methyluric acid intermediate (24):

The nitrosopyrimidine **19** (270 mg, 1.06 mmol) was dissolved in ethanol (20 mL) with warming and palladium on carbon (75 mg, 10%) was added under argon. Hydrogenation was performed at room temperature and 15 psi for 2 h, filtered to remove catalyst and evaporated to dryness. The residue was combined with urea (600 mg, 10 mmol) and heated neat on the hot plate with stirring. The temperature reached 140°C which produced a clear melt and was maintained for about 10 min. with additional urea added (1 g). Upon cooling the melt solidified and was dissolved in n NaOH (25 mL) and boiled with decolorizing carbon for 10 min., filtered and acidified to pH 3-4 while hot. The resulting precipitate was collected after cooling and washed with water and dried to yield 160 mg (57%) of **24** as an off-white solid with the following characteristics: mp >290°C dec. ¹H NMR (500 MHz, DMSO-d₆) δ 11.80 and 10.73 (2s, 2H, N-7 H, N-9 H), 3.78 (t, 2H, N-CH₂), 3.30 (s, 3H, N-CH₃, under H₂O signal), 1.48 (m, 2H, 2'CH₂), 1.24 (m, 6H, 3', 4', 5' CH₂), 0.85 (t, 3H, CH₃). Analysis: C₁₂H₁₈N₄O₃ (C, H, N; Table 2).

Synthesis of 3-Methyl-8-thiouric acid (25) intermediate:

The pyrimidinediamine **33** (100 mg, 0.63 mmol) was combined with potassium ethyl xanthate (810 mg, 5 mmol) and DMF (10 mL) and heated at 100°C. The suspension became green almost immediately and reaction was complete after 30 min. by TLC. After a total reaction time of 1 h, the mixture was cooled, filtered and washed with Et₂O, dried to yield an off-white solid (310 mg) which presumably contained the unreacted potassium ethyl xanthate and the potassium salt of the desired

product. The solid was suspended in water (5 mL) and heated to dissolve. Glacial acetic acid was added to pH 5 and a vigorous effervescence was noted. A white solid formed which was filtered warm and washed with water, then ethanol and dried to yield 99 mg (79%) of the title compound. ¹H NMR (DMSO-d₆) δ 13.40, 12.92 and
5 11.80 (3br s, 3H, NHs), 3.28 (s, 3H, CH₃). Analysis: C₆H₆N₄O₂S (C, H, N; Table 2).

Synthesis of 3-n-Propylxanthine (29) intermediate:

The pyrimidinediamine 28 (750 mg) was combined with diethoxymethyl acetate (7 mL) and heated at 80°C for 2 h. The mixture was evaporated to dryness and water (5 mL) was added and the mixture heated for 20 min. to near boiling. The
10 resulting solution was then allowed to evaporate slowly to yield off-white crystals. Yield 680 mg (86%); mp 282-284°C, Lit.¹⁵ 291-292°C.

EXAMPLE XIV

SYNTHESIS OF COMPOUNDS 36-39, 41 AND 43 AND INTERMEDIATES

General Procedure for Ring Closure of Pyrimidine-diamines to Pteridines:

15 The orthodiamine 28 or 33 (2 mmol) was suspended in water (20 mL) and heated to above 70°C before a solution of glyoxal-sodium bisulfite addition product (10 mmol in 25 mL water) was added with stirring. The pale green suspension slowly became light amber and clear. After heating 5 min TLC indicated reaction was complete. The mixture was cooled and extracted with ethyl acetate (5 x 40 mL), dried
20 over MgSO₄ and evaporated to yield the 1-methyl (34) or 1-n-propylpteridine (35) in

71 and 78 %, respectively. ¹H NMR showed the appearance of two aromatic signals at about 8.74 and 8.55 as doublets (J = 2.5 Hz) for both compounds.

Synthesis of 6,7-Diethyl-1-methylpteridine-2,4-dione (40) intermediate:

Compound 33 (200 mg, 1.27 mmol) was suspended in acetonitrile (5 mL) and 3,4-hexanedione (185 μ L, 1.52 mmol) was added. The mixture was heated at 70°C for 15 min with minimal product formation due to insolubility of 33. Therefore DMF (3 mL) and water (3 mL) were added and the temperature was raised to 100°C. After 90 min total reaction time the mixture was cooled and poured into water (100 mL) and extracted with ethyl acetate (3 x 75 mL). The organic layer was dried over MgSO₄ and evaporated to provide the colorless crystalline product. Yield 240 mg (81%); mp 218-222°C; ¹H NMR (DMSO-d₆) δ 11.78 (br s, 1H, NH), 3.46 (s, 3H, NCH₃), 2.95 and 2.93 (2q, 4H, 2CH₂ of ethyls), 1.28 and 1.23 (2t, 6H, 2CH₃ of ethyls). Analysis: C₁₁H₁₄N₄O₂ (C, H, N; Table 2).

Synthesis of 1-Methyl-6-phenylpteridine-2,4-dione (42) intermediate:

The nitrosopyrimidine 32 (220 mg, 1.28 mmol) was mixed thoroughly with phenethyl amine hydrochloride (1.5 g, 9.5 mmol) and heated in an open beaker on the hot plate. After a few minutes at about 160°C the purple reaction mixture fused to a brown paste. TLC indicated many products so sulfolane (1 mL) was added and heat was continued for 15 min. The reaction mixture was heated in water (10 mL) and then diluted 50 mL in water and extracted with ethyl acetate (2 x 50 mL), the organic layer dried over MgSO₄ and then concentrated. The residue was flash

chromatographed on silica gel using 4% MeOH in CH_2Cl_2 . Yield 75 mg (23%) of 42 as a pale yellow-orange solid. mp $>307^\circ\text{C}$ dec.; ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 11.95 (br s, 1H, NH), 9.37 (s, 1H, C-7 H), 8.17 (m, 2H, 2',6' phenyl), 7.55 (m, 3H, 3',4',5' phenyl), 3.51 (s, 3H, NCH_3). Anal. $\text{C}_{13}\text{H}_{10}\text{N}_4\text{O}_2$ (C, H, N).

EXAMPLE XV

SYNTHESIS OF COMPOUNDS 44, 47 AND 48

General Method for Ring Closure of Pyrimidines to Thiadiazolo-pyrimidines (Compounds 44-46):

5 The orthodiamine 23, 27, or 32 (2.3 mmol) was suspended in dry acetonitrile (5 mL) and dry pyridine (1.5 mL) was added. Thionyl chloride (1 mL, 13.7 mmol) was added quickly and the mixture, which became clear and darkened, was heated at 60°C for 10 min. The mixture was then cooled and poured into n HCl (40 mL) with stirring. The resulting yellow solution was extracted with ethyl acetate
10 (3 x 40 mL), dried over MgSO₄ and evaporated to yield a pale yellow solid which was triturated with ether. Yield 65-74%.

Alkylation of these intermediates yielded the disubstituted products 47 and 48.

EXAMPLE XVI

SYNTHESIS OF COMPOUNDS 50 AND 52

Ethyl 4-[(2-methylamino)benzoyl]aminobutanoate (51):

15 A mixture of N-methylisatoic anhydride (3.5 g, 19.8 mmol) was combined with 4-aminobutyric acid (2.5 g, 24.3 mmol) in dry DMF (50 mL) and heated at 100°C for 2 h. TLC indicated reaction to be complete and the DMF was removed in
20 vacuo. The residue was used directly for esterification which was accomplished by dissolving the residue in 100% ethanol (50 mL) and adding chlorotrimethyl silane (2.5 mL, 20 mmol). The mixture was heated at 65°C for 6 h and then evaporated to yield

a brown syrup. Crude yield 87% from isatoic anhydride. A small sample was purified for characterization and biological testing by preparative TLC using 7% MeOH in CH₂Cl₂. The remainder of the material was used directly for preparation of compound 52. Analysis: C₁₄H₂₀N₂O₃ (C, H, N; Table 2).

5 **Ethyl 1-Methyl-1,4-dihydro-2,4-dioxo-3(2*H*)-quinazolinebutanoate (52):**

The residue from 51 was combined with ethyl chloroformate (10 mL) and heated at 90°C for 1 h. The mixture was cooled and poured into saturated aqueous sodium bicarbonate (50 mL) with stirring and after 10 min extracted with ethyl acetate
10 (2 x 75 mL). The organic layer was dried over MgSO₄ and evaporated to yield a brown syrup. The crude product was flash chromatographed on silica using 3% MeOH in CH₂Cl₂ to yield g (%) of 52 as a thick oil. ¹H NMR (500 MHz, DMSO-d₆) δ 7.27-7.42 (2m, 4H, C-5,6,7,8), 4.04 (t, 2H, CH₂ of ethyl), 3.88 (m, 2H, NCH₂), 3.11 (s, 3H, NCH₃), 2.33 (t, 2H, 2'CH₂), 1.71 (m, 2H, 3'CH₂). Analysis: C₁₅H₁₈N₂O₄ (C,
15 H, N).

EXAMPLE XVII

SYNTHESIS OF HIGHLY WATER SOLUBLE DERIVATIVES

Acid Hydrolysis of Compound 53.

Compound 53 (1.63 g, 5.1 mmol) was combined with n HCl (30 mL) and
20 was heated at boiling for 75 min. The mixture was cooled and the resulting solid which formed was filtered, washed with cold water, and dried to yield 1.5 g

(quantitative conversion) of the free carboxylic acid of suitable purity for esterification (as judged by TLC).


Esterification to form morpholinoethyl ester Compound 54.

The carboxylic acid obtained above in this Example (1.0 g, 3.4 mmol) was dissolved in dichloromethane (25 mL) with warming and then thionyl chloride (1 mL, 13.7 mmol) was added followed by 3 drops of DMF. After a few minutes a white solid precipitated (acid chloride intermediate), but the reaction mixture was allowed to stir at room temperature overnight. The mixture was evaporated to remove excess thionyl chloride and the residue was suspended in dry acetonitrile (25 mL). To this mixture was added morpholinoethanol (1.24 mL, 10.2 mmol) and heated at 80°C for 5 minutes. The solid suspension became clear almost immediately and after cooling, the reaction mixture was evaporated onto silica gel and loaded on a flash silica gel column and eluted with methanol - dichloromethane 5/95. The fractions of pure product were pooled and evaporated to yield an off-white residue which was dissolved in isopropyl alcohol (5 mL) and concentrated HCl (1 mL) was added. Upon concentration in vacuo, an off-white solid formed. Yield after drying 0.93 g (62%), mp 90°C. Proton NMR confirms structure assignment as the morpholino ethyl ester of Compound 53: ¹H NMR (DMSO-d₆, 500 MHz) δ 7.56 and 7.15 (2s, 2H, C-5 and C-8 aromatics), 7.35 and 6.55 (2d, 2H, C-3 and C-4 aromatics), 4.37 (m, 2H, C-1' of ester), 3.92 (m, 2H, C-4 of acyl), 3.86 and 3.84 (2s, 6H, OMe's), 3.95, 3.45, 3.10 (3m, 8H, morpholino and C-2'), 2.38 (t, 2H, C-2 of acyl), 1.92 (m, 2H, C-3 of acyl).

The invention having been fully described, modifications thereof may be apparent to those of ordinary skill in the art. Such modifications are within the scope of the invention as defined by the appended claims.

CLAIMS

1. A compound for antagonizing activation of c-Jun kinase and p38 comprised of a first constituent having the formula:



wherein:

- 5 R_1 is a terminally substituted normal alkyl having from 1 to 7 carbon atoms, a terminally substituted alkenyl having from 2 to 7 carbon atoms, a terminally substituted ether having from 2 to 6 carbon atoms, a terminally substituted secondary amine having from 2 to 6 carbon atoms, or substituted aryl having less than 8 carbons, where said terminal group is
- 10 NH_2 , substituted amino, acyloxy, SO_3H , PO_4H_2 , $NNO(OH)$, SO_2NH_2 , $PO(OH)NH_2$, SO_2R or $COOR$, where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl, benzyl, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is NH_2 or a substituted amino where the substituents

on the amino have 1 to 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom;

Z is C, CH, or N;

5 R₂ is an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms when Z is C, R₂ is a halogen, NO, amino, or substituted amino when Z is CH, or R₂ is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms when Z is N;

10 A is CO when Z is N, or CR₅ when Z is C or CH, where R₅ is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, aryl or aralkyl having less than 7 carbon atoms, OH, or an O-alkyl having from 1 to 5 carbon atoms and there is a double bond between Z and A when Z is C and a single bond between Z and A when Z is CH;

15 Y₁ is N, NR₆ or CR₆, where R₆ is H, NO, an amino, a substituted amino, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms;

Y₂ is N or CH; and

20 X is S when Y₁ and Y₂ are N, CR₇ when Y₁ is NR₆ where R₇ is H, OH, SH, Br, Cl, or I, or =C(R₃)-C(R₄)= when Y₁ is N, CH or CR₆ and Y₂ is N or CH, where each of R₃ and R₄, independently, is H, an alkyl, a cyclic alkyl,


a heterocyclic alkyl, alkenyl, aryl, alkylcarboxyl, or aralkyl having less than 7 carbon atoms, SH, OH or an O-alkyl having from 1 to 5 carbon atoms,

or a salt thereof; and,

- 5 a second constituent consisting of a pyridinylimidazole conjugated to the first constituent;

wherein the first constituent selectively antagonizes c-Jun kinase activation and the second constituent selectively antagonizes p38 activation.

2. The compound according to Claim 1, wherein the first constituent has the formula:



wherein

5 R₁ is a terminally substituted normal alkyl having from 1 to 7 carbon atoms, a terminally substituted alkenyl having from 2 to 7 carbon atoms, a terminally substituted ether having from 2 to 6 carbon atoms, a terminally substituted secondary amine having from 2 to 6 carbon atoms, or substituted aryl having less than 8 carbons, where said terminal group is
 10 NH₂, substituted amino, acyloxy, SO₃H, PO₄H₂, NNO(OH), SO₂NH₂, PO(OH)NH₂, SO₂R or COOR where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl, benzyl, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is NH₂ or a substituted amino where the substituents
 15 on the amino have 1 to 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom;

R_2 is an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms;

each of Y_1 and Y_2 , independently, is N or CH;


5 each of R_3 and R_4 , independently, is H, SH, OH, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, aryl, alkylcarboxyl, or aralkyl having less than 7 carbon atoms, or OH or an O-alkyl having from 1 to 5 carbon atoms;

R_5 , when present, is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, aryl or aralkyl having less than 7 carbon atoms, OH, or an O-alkyl having from 1 to 5 carbon atoms; and


10 R_6 is NO, an amino, or a substituted amino;

or a salt thereof.

3. The compound of Claim 2 wherein the first constituent has the formula:



or



wherein

5 R₁ is a terminally substituted normal alkyl having from 1 to 7 carbon atoms, a terminally substituted alkenyl having from 2 to 7 carbon atoms, a terminally substituted ether having from 2 to 6 carbon atoms, a terminally substituted secondary amine having from 2 to 6 carbon atoms, or substituted aryl having less than 8 carbons, where said terminal group is NH₂, substituted amino, acyloxy, SO₃H, PO₄H₂, NNO(OH), SO₂NH₂, PO(OH)NH₂, SO₂R or COOR where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl, 10 benzyl, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is NH₂ or a substituted amino where the substituents on the amino have 1 to 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom;

15 R₂, when present, is an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms; and

each of R₃ and R₄, independently, is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, aryl, alkylcarboxyl, or aralkyl having less than 7 carbon atoms, or OH or an O-alkyl having from 1 to 5 carbon atoms; or a salt thereof.


4. The compound of Claim 3, wherein the first constituent has the following substituents:

5 R_1 is a terminally substituted normal alkyl having from 1 to 7 carbon atoms, where said terminal group and is NH_2 , substituted amino, or $COOR$, where R is H, an alkyl having from 1 to 4 carbon atoms, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is a substituted amino where the substituents on the amino have 1 to 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom;

10 R_2 , when present, is an alkyl having less than 7 carbon atoms; and each of R_3 and R_4 , independently, is H, an alkyl, aryl, alkylcarboxyl, or aralkyl having less than 7 carbon atoms, or an O-alkyl having from 1 to 5 carbon atoms;

or a salt thereof.

5. The compound of Claim 4 wherein the first constituent has the formula:



wherein

5 R_1 is a terminally substituted normal alkyl having from 1 to 7 carbon atoms where said terminal group is COOR, where R is H, an alkyl having from 1 to 4 carbon atoms, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is a morpholino group; and

each of R_3 and R_4 , independently, is H, an alkyl having less than 7 carbon atoms, or an O-alkyl having from 1 to 5 carbon atoms;

10 or a salt thereof.

6. The compound of Claim 5, wherein the first constituent has the following substituents:

R_1 is a terminally substituted normal alkyl having from 1 to 4 carbon atoms where said terminal group is COOR where R is an N-morpholinoalkyl

5 group; and,


each of R_3 and R_4 , independently, is an O-alkyl having from 1 to 5 carbon atoms;

or a salt thereof.

7. The compound of Claim 6, wherein R_1 is $-(CH_2)_3COOR$ where R is an N-morpholinoethyl group, and R_3 and R_4 each are methoxy groups, or a salt thereof.

10

8. The compound of Claim 4, wherein the first constituent has the formula:



wherein

R₁ is a terminally substituted normal alkyl having from 1 to 7 carbon atoms
5 where said terminal group is COOR, where R is H, an alkyl having from 1 to 4 carbon atoms, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is a morpholino group;

R₂ is an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms; and

10 each of R₃ and R₄, independently, is H, OH, SH, an alkyl, aryl, or aralkyl having less than 7 carbon atoms, or an O-alkyl having from 1 to 5 carbon atoms;

or a salt thereof.

9. The compound of Claim 8, wherein the first constituent has the following
15 substituents:

R_1 is a terminally substituted normal alkyl having from 1 to 4 carbon atoms where said terminal group is COOR where R is an N-morpholinoalkyl group;

R_2 is an alkyl having less than 7 carbon atoms; and

5 each of R_3 and R_4 , independently, is H, an alkyl, aryl, or alkylcarboxyl having less than 7 carbon atoms, or an O-alkyl having from 1 to 5 carbon atoms;

or a salt thereof.

10. The compound of Claim 9, wherein the first constituent has the following
10 substituents:

R_1 is $-(CH_2)_3COOR$ where R is an N-morpholinoethyl group;

R_2 is an alkyl having less than 4 carbon atoms; and

each of R_3 and R_4 , independently, is H, an alkyl having 1 to 3 carbon atoms, phenyl, or methylcarboxyl;

15 or a salt thereof.


11. The compound of Claim 10, wherein the first constituent has the following substituents:

R_2 is $-(CH_2)_2CH_3$; and,

R_3 and R_4 each are H;

5 or a salt thereof.

12. The compound according to Claim 1 wherein the first constituent has the formula:



wherein

10 R_1 is a terminally substituted normal alkyl having from 1 to 7 carbon atoms, a terminally substituted alkenyl having from 2 to 7 carbon atoms, a terminally substituted ether having from 2 to 6 carbon atoms, a terminally substituted secondary amine having from 2 to 6 carbon atoms, or substituted aryl having less than 8 carbons, where said terminal group is NH_2 , substituted amino, acyloxy, SO_3H , PO_4H_2 , $NNO(OH)$, SO_2NH_2 ,

PO(OH)NH₂, SO₂R or COOR, where R is H, an alkyl having from 1 to 4 carbon atoms, an alkenyl having from 1 to 4 carbon atoms, tetrazolyl, benzyl, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is NH₂ or a substituted amino where the substituents on the amino have 1 to 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom;

R₂ is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms;

Y₂ is N or CH;

R₆ is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms; and

R₇ is H, OH, SH, Br, Cl, or I;

or a salt thereof; and,

a second constituent consisting of a pyridinylimidazole;

wherein the first constituent selectively antagonizes c-Jun kinase activation and the second constituent selectively antagonizes p38 activation.

13. The compound of Claim 12, wherein the first constituent has the following substituents:

R_1 is a terminally substituted normal alkyl having from 1 to 7 carbon atoms, where said terminal group is NH_2 , substituted amino, or $COOR$, where R is H, an alkyl having from 1 to 4 carbon atoms, or an alkylamino, where the alkyl group has from 1 to 4 carbon atoms and the amino is NH_2 or a substituted amino where the substituents on the amino have 1 to 6 carbon atoms, one of which can be replaced by an oxygen atom or nitrogen atom;

R_2 is H, or an alkyl having less than 7 carbon atoms;

Y_2 is N or CH;

R_6 is H, or an alkyl having less than 7 carbon atoms; and

R_7 is H, OH, SH, or Br,

or a salt thereof.

14. The compound of Claim 13, wherein R_1 is $(CH_2)_3COOR$ and R is an N-morpholinoethyl group, or a salt thereof.

15. The compound of Claim 1, wherein R_1 is $(CH_2)_3COOR$ and R is an N-morpholinoethyl group, or a salt thereof.

16. The compound according to Claim 1, wherein the second constituent is a cytokine suppressive anti-inflammatory drug.

17. The compound according to Claim 16, wherein the cytokine suppressive anti-inflammatory drug is selected from the group of such drugs comprising 1-[3-(4-morpholinyl)propyl]-4-(4-fluorophenyl)-5-(4-pyridyl)imidazole, 2-(4-methylsulfinylphenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 2-(4-methylthiophenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 4-(4-fluorophenyl)-2-(4-methylsulfinyl)-5-(4-pyridyl)imidazole and 5-(4-pyridyl)-6-(4-fluorophenyl)-2,3-dihydroimidazole(2,1-b)thiazole.


18. The compound according to Claim 1, wherein the first and second constituents are conjugated to one another by a bond which is severable *in vivo*.

19. The compound according to Claim 18, wherein the bond is an amide, ester or azo bond.


20. The compound according to Claim 19, wherein the bond is a tertiary N-acyloxymethyl amide bond.

21. The compound according to Claim 1 mixed in a colloid.
22. The compound according to Claim 21, wherein the colloid is a liposome.
23. A pharmaceutical composition comprising the compound of Claim 1 and a pharmaceutically acceptable carrier.

24. A compound for antagonizing activation of c-Jun kinase and p38 having the formula:



or



wherein

5

n is any number of carbon atoms from 1 to 7, O or N;

R_1 , if present, is H, an alkyl, a cyclic alkyl, a heterocyclic alkyl, alkenyl, or aralkyl having less than 7 carbon atoms;

R_6 and R_7 are H, OH or OR_1 , in any combination;


Z is N or C;

10


X, where Z is C, is H, halogen, N_3 , NO, NH_2 , NHR_1 , $N(R_1)_2$ or COR_1 ; and,

A is H, halogen, N_3 , NO, NH_2 , NHR_1 , $N(R_1)_2$ or COR_1 .

25. A compound according to Claim 24 having the formula:




26. A compound according to Claim 24 having the formula:



27. A pharmaceutically acceptable composition comprising the compound according to Claim 24 mixed in a colloid.

28. The compound according to Claim 27, wherein the colloid is a liposome.
29. A pharmaceutical composition comprising the compound of Claim 24 and a pharmaceutically acceptable carrier.
30. A compound comprised of a first constituent having the formula:



5 wherein

R₁ is (CH₂)COO-Et;

R₂ is CH₃;

R₃ is CH₃; and,

Z is N;

further comprised of a second constituent conjugated to the first constituent and consisting of a pyridinylimidazole which selectively antagonizes p38 activation.


31. The compound according to Claim 30, wherein the pyridinylimidazole is a
5 cytokine suppressive anti-inflammatory drug.

32. The compound according to Claim 31, wherein the cytokine suppressive anti-inflammatory drug is selected from the group of such drugs comprising 1-[3-(4-morpholinyl)propyl]-4-(4-fluorophenyl)-5-(4-pyridyl)imidazole, 2-(4-methylsulfinylphenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 2-(4-methylthiophenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 4-(4-fluorophenyl)-2-(4-methylsulfinyl)-5-(4-pyridyl)imidazole and 5-(4-pyridyl)-6-(4-fluorophenyl)-2,3-dihydroimidazole(2,1-b)thiazole.

10

33. The compound according to Claim 30, wherein the first and second constituents are conjugated to one another by a bond which is severable *in vivo*.

34. The compound according to Claim 33, wherein the bond is an amide, ester or azo bond.
35. The compound according to Claim 34, wherein the bond is a tertiary N acyloxymethyl amide bond.
- 5 36. The compound according to Claim 30, wherein the first and second constituents are mixed in a colloid.
37. The compound according to Claim 36, wherein the colloid is a liposome.
38. A pharmaceutical composition comprising the compound of Claim 30 and a pharmaceutically acceptable carrier.
- 10 39. A compound comprised of a first constituent having the formula:



wherein

R_1 is $(CH_2)_3$ COO-Et;

R_2 is n-propyl;

R_4 is H; and

5 R_5 is H;

further comprised of a second constituent conjugated to the first constituent and consisting of a pyridinylimidazole which selectively antagonizes p38 activation.

40. The compound according to Claim 39, wherein the pyridinylimidazole is a
10 cytokine suppressive anti-inflammatory drug.

41. The compound according to Claim 40, wherein the cytokine suppressive anti-inflammatory drug is selected from the group of such drugs comprising 1-[3-(4-

- morpholinyl)propyl]-4-(4-fluorophenyl)-5-(4-pyridyl)imidazole, 2-(4-methylsulfinylphenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 2-(4-methylthiophenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 4-(4-fluorophenyl)-2-(4-methylsulfinyl)-5-(4-pyridyl)imidazole and 5-(4-pyridyl)-6-(4-fluorophenyl)-2,3-dihydroimidazole(2,1-b)thiazole.
- 5

42. The compound according to Claim 39, wherein the first and second constituents are conjugated to one another by a bond which is severable *in vivo*.

43. The compound according to Claim 42, wherein the bond is an amide, ester or azo bond.


- 10 44. The compound according to Claim 43, wherein the bond is a tertiary N-acyloxymethyl amide bond.

45. The compound according to Claim 39 mixed in a colloid.

46. The compound according to Claim 45, wherein the colloid is a liposome.

47. A pharmaceutical composition comprising the compound of Claim 39 and a pharmaceutically acceptable carrier.

48. A compound comprised of a first constituent having the formula:




53

1) NaOH or HCl, heat

2) SOCl₂

3) morpholinoethanol

4) HCl



54

further comprised of a second constituent conjugated to the first constituent and consisting of a pyridinylimidazole which selectively antagonizes p38 activation.

49. The compound according to Claim 48, wherein the pyridinylimidazole is a
5 cytokine suppressive anti-inflammatory drug.

50. The compound according to Claim 49, wherein the cytokine suppressive anti-inflammatory drug is selected from the group of such drugs comprising 1-[3-(4-morpholinyl)propyl]-4-(4-fluorophenyl)-5-(4-pyridyl)imidazole, 2-(4-methylsulfinylphenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 2-(4-methylthiophenyl)-3-(4-pyridyl)-6,7-dihydro-[5H]-pyrrolo[1,2-a]imidazole, 4-(4-fluorophenyl)-2-(4-methylsulfinyl)-5-(4-pyridyl)imidazole and 5-(4-pyridyl)-6-(4-fluorophenyl)-2,3-dihydroimidazole(2,1-b)thiazole.

51. The compound according to Claim 48, wherein the first and second constituents are conjugated to one another by a bond which is severable *in vivo*.

- 10 52. The compound according to Claim 51, wherein the bond is an amide, ester or azo bond.

53. The compound according to Claim 52, wherein the bond is a tertiary N-acyloxymethyl amide bond.

54. The compound according to Claim 48 mixed in a colloid.

55. The compound according to Claim 54, wherein the colloid is a liposome.
56. A pharmaceutical composition comprising the compound of Claim 48 and a pharmaceutically acceptable carrier.
57. A method for treating a vertebrate host for an inflammatory condition
5 comprising administering the compound of Claim 1 to the host.
58. The method according to Claim 57 wherein the inflammatory condition is arthritis, spondylitis, bone resorption, sepsis, septic shock, endotoxic shock, inflammatory bowel disease, asthma, bronchitis, chronic pulmonary inflammatory disease, silicosis, reperfusion injury, graft versus host reactions, allograft rejection,
10 fever, viral and bacterial infection, inflammation of the joints, psoriasis and eczema, radiation dermatitis, multiple sclerosis, cell senescence and apoptosis.
59. A method for treating a vertebrate host for an inflammatory condition comprising administering the compound of Claim 24 to the host.

60. The method according to Claim 59 wherein the inflammatory condition is arthritis, spondylitis, bone resorption, sepsis, septic shock, endotoxic shock, inflammatory bowel disease, asthma, bronchitis, chronic pulmonary inflammatory disease, silicosis, reperfusion injury, graft versus host reactions, allograft rejection, fever, viral and bacterial infection, inflammation of the joints, psoriasis and eczema, radiation dermatitis, multiple sclerosis, cell senescence and apoptosis.

61. A method for treating a vertebrate host for an inflammatory condition comprising administering the compound of Claim 30 to the host.

62. The method according to Claim 61 wherein the inflammatory condition is arthritis, spondylitis, bone resorption, sepsis, septic shock, endotoxic shock, inflammatory bowel disease, asthma, bronchitis, chronic pulmonary inflammatory disease, silicosis, reperfusion injury, graft versus host reactions, allograft rejection, fever, viral and bacterial infection, inflammation of the joints, psoriasis and eczema, radiation dermatitis, multiple sclerosis, cell senescence and apoptosis.

63. A method for treating a vertebrate host for an inflammatory condition comprising administering the compound of Claim 39 to the host.

64. The method according to Claim 63 wherein the inflammatory condition is arthritis, spondylitis, bone resorption, sepsis, septic shock, endotoxic shock, inflammatory bowel disease, asthma, bronchitis, chronic pulmonary inflammatory disease, silicosis, reperfusion injury, graft versus host reactions, allograft rejection, 5 fever, viral and bacterial infection, inflammation of the joints, psoriasis and eczema, radiation dermatitis, multiple sclerosis, cell senescence and apoptosis.

65. A method for treating a vertebrate host for an inflammatory condition comprising administering the compound of Claim 48 to the host.

66. The method according to Claim 65 wherein the inflammatory condition is 10 arthritis, spondylitis, bone resorption, sepsis, septic shock, endotoxic shock, inflammatory bowel disease, asthma, bronchitis, chronic pulmonary inflammatory disease, silicosis, reperfusion injury, graft versus host reactions, allograft rejection, fever, viral and bacterial infection, inflammation of the joints, psoriasis and eczema, radiation dermatitis, multiple sclerosis, cell senescence and apoptosis.

1/14




FIG. 1





FIG. 2

2/14

FIG. 3



3/14

FIG. 4(A)




FIG. 4(B)





FIG. 4(C)



4/14

FIG. 4(D)




FIG. 4(E)





FIG. 4(F)



5/14

FIG. 4(G)




FIG. 5(b)





FIG. 5(a)



7/14

FIG. 6





FIG. 9



8/14




FIG. 7

9/14




FIG. 8

10/14

FIG. 10






FIG. 11



11/14


FIG. 12




12/14

S45,928-3


3-NITROIXOCARBOSTYRIL

 $C_9H_6N_2O_3$ 


S76,334-9

N2DIMMETHYL-3-(3-FLUOROPHENYL)-1-OXO-1234
TETRAHYDRO-4-ISOQUINOLINECARBOXAMIDE $C_{18}H_{17}FN_2O_2$ 

S93,63C-8


3-AMINO-CYANO-7-NITRO-2-PHENYL-1(2H)-
ISOQUINOLONE $C_{15}H_{10}N_4O_3$ 

S75,469-8


2-METHYL-3-(2-OXOPROPYL)-1(2H)-ISOQUINOLINE
CNE $C_{13}H_{13}NO_2$ 

S88,370-0

2-(2-HYDROXYETHYL)-1,4-(2H,3H)-ISOQUINOLONE

 $C_{11}H_{11}NO_3$ 

21,493-0


(H)-1234 TETRAHYDRO-3-ISOQUINOLINE
CARBOXYLIC ACID HYDROCHLORIDE $C_{10}H_{11}NO_2$ 

*HC


FIG. 13(A)

13/14

36,895-4
3-HYDROXYISOQUINOLINE, 99%
 C_9H_7NO



S94,009-7
3-AMINO-4-CYANO-2-HEXYL-7-NITRO-1(2H)-
ISOQUINOLONE
 $C_{16}H_{18}N_4O_3$



S75,699-7
2-BROMO-4-NITRO-6(5H)-PHENANTHRIDINONE
 $C_{13}H_7BrN_2O_3$






FIG. 13(B)


S52,626-6
6,7-DIMETHOXY-1(2H)-ISOQUINOLONE
 $C_{11}H_{11}NO_3$



S84,100-5
4-ISOQUINOLINECARBOXYLIC ACID
 $C_{10}H_7NO_2$




15,013-4
1-ISOQUINOLINECARBOXYLIC ACID, 99%
 $C_{10}H_7NO_2$




14/14

S76,087-0

2-(2-(3,4-DIMETHOXYPHENYL)ETHYL)ISOQUINOLINE-1,3
(2H,4H)-DIONE $C_{19}H_{19}NO_4$ 

S76,473-6

1,2-DIHYDRO-2-METHYL-1-OXO-3-ISOQUINOLINES
UTYRIC ACID $C_{14}H_{15}NO_3$ 

29,963-4

5(5H)-PHENANTHRIDINONE



 $C_{13}H_9NO$ 

FIG. 13(C)


N165-8

1,8-NAPHTHALIMIDE, 99%

 $C_{12}H_7NO_2$ 

33,854-0

3-ISOQUINOLINECARBOXYLIC ACID HYDRATE, 99%

 $C_{10}H_7NO_2$ 

PCT/US 99/14320

CLASSIFICATION OF SUBJECT MATTER				
IPC 7	C07D475/02	C07D473/04	C07D513/04	C07D239/96 C07D217/24
	C07D401/14	A61K31/505	A61K31/52	/(C07D513/04, 285:00,
	239:00)			

According to International Patent Classification (IPC) or to both national classification and IPC

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 20710 A (UNIV CALIFORNIA) 11 July 1996 (1996-07-11) cited in the application claims ---	1-66
Y	WO 94 22449 A (CELL THERAPEUTICS INC) 13 October 1994 (1994-10-13) claims ---	1-66
A	WO 94 22863 A (CELL THERAPEUTICS INC) 13 October 1994 (1994-10-13) claims ---	1-66
A	WO 92 21344 A (HUTCHINSON FRED CANCER RES) 10 December 1992 (1992-12-10) claims ---	1-66

	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

"&" document member of the same patent family

15 October 1999

22/10/1999

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Chouly, J

INTERNATIONAL SEARCH REPORT

Inter- national Application No

PCT/US 99/14320

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	<p>WO 98 52948 A (UNIV CALIFORNIA) 26 November 1998 (1998-11-26) cited in the application claims 1-66</p> <p>-----</p>	1-66

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/14320

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 57-66
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 57-66
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/14320

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9620710	A	11-07-1996	US 5843943 A AU 4601996 A CA 2208580 A EP 0801568 A JP 11502193 T	01-12-1998 24-07-1996 11-07-1996 22-10-1997 23-02-1998
WO 9422449	A	13-10-1994	US 5670506 A AU 6623894 A EP 0714302 A	23-09-1997 24-10-1994 05-06-1996
WO 9422863	A	13-10-1994	US 5837703 A US 5470878 A AU 695674 B AU 6553894 A AU 9051898 A CA 2159640 A CH 686830 A CN 1122600 A, B EP 0719267 A JP 8508981 T NZ 265178 A US 5777117 A US 5824677 A US 5801181 A US 5750575 A ZA 9402317 A US 5641783 A	17-11-1998 28-11-1995 20-08-1998 24-10-1994 14-01-1999 13-10-1994 15-07-1996 15-05-1996 03-07-1996 24-09-1996 27-04-1995 07-07-1998 20-10-1998 01-09-1998 12-05-1998 10-02-1995 24-06-1997
WO 9221344	A	10-12-1992	AU 664189 B AU 2247592 A CA 2109796 A EP 0573617 A JP 7500085 T US 5856115 A US 5585380 A	09-11-1995 08-01-1993 10-12-1992 15-12-1993 05-01-1995 05-01-1999 17-12-1996
WO 9852948	A	26-11-1998	AU 7489698 A	11-12-1998